

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure

Deputy 250WG

Herbicide

ACTIVE CONSTITUENT: 250 g/kg BUTROXYDIM

GROUP	1	HERBICIDE
-------	---	-----------

For the control of certain grasses in a range of broadacre crops as per the directions for use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT

NET CONTENTS: 1 kg - 50kg

CropSure Pty Ltd

18 Raymond Road

Laverton North

Victoria 3026

ABN: 17 643 829 190

ACN: 643 829 190

Phone: 03 9931 2200

www.cropsure.com

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store in prolonged periods of direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling dust. When preparing product for use wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126; New Zealand 0800 764 766

SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet, which is available from the supplier

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

24HR Medical Emergency Information:
Australia +61 1800 951 288
New Zealand +64 800 700 112

APVMA APPROVAL NO: 92222/134877

BATCH NO:

DATE OF MANUFACTURE:

Harmful if swallowed. Causes skin irritation. Suspected of causing cancer. Suspected of damaging fertility. Suspected of damaging the unborn child. Very toxic to aquatic life with long lasting effects. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves, protective clothing and eye or face protection. Brush off loose particles from skin. Take off contaminated clothing and wash before reuse. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN: Wash with plenty of soap and water. If exposed or concerned: Get medical advice. If skin irritation occurs: Get medical advice. In case of fire: Use carbon dioxide, dry chemical, foam, water fog, to extinguish. Store locked up. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure

Deputy 250WG

Herbicide

ACTIVE CONSTITUENT: 250 g/kg BUTROXYDIM

GROUP	1	HERBICIDE
-------	---	-----------

For the control of certain grasses in a range of broadacre crops as per the directions for use.

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

APVMA APPROVAL NO: 92222/134877

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

DIRECTIONS FOR USE**Restraints:**

DO NOT treat weeds that are not actively growing or are growing under stress. Under such circumstances the biological processes of the weeds slow down and CropSure Deputy 250WG Herbicide transport in the weed can be drastically reduced, resulting in an incomplete kill or suppression only of the weeds.

DO NOT apply to annual grass weeds after mid-tillering.

DO NOT apply at flowering stage of weeds.

DO NOT apply under conditions of prolonged high or very low temperature (frosts), moisture stress (waterlogging or drought), low humidity, poor fertility or stress from previous herbicide application.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

CROP	WEEDS CONTROLLED	WEED STAGE	STAT E	RATE /ha	CRITICAL COMMENTS
Chickpeas Faba beans Field peas Lentils Linseed Lucerne Lupins Vetches (<i>Vicia</i> spp.) White and Subterranean Clover and Medic pastures	Annual ryegrass (<i>Lolium rigidum</i>) Barley grass (<i>Hordeum</i> spp.) Wild oats (<i>Avena</i> spp.)	2 leaf to early tillering Zadoks 12-23	All States	80 to 180 g *	Use the lower rates for younger weeds growing actively under ideal conditions. Use the higher rates for weeds that are predominantly at early tillering, or where denser populations are present or under less than ideal growing conditions or where some weed resistance to Group A herbicides is suspected. Because of the extreme variation of cross resistance present in annual ryegrass populations there is no guarantee that CropSure Deputy 250WG Herbicide will always provide consistently high levels of control but trials have shown that higher rates may be expected to perform better than lower rates. For all crops apply in not less than 50 L water/ha.
	Annual ryegrass (<i>Lolium rigidum</i>) together with any of the following grasses: Annual phalaris (<i>Phalaris</i> spp.), Barley grass (<i>Hordeum</i> spp.), Brome grass (<i>Bromus</i> spp.) Volunteer cereals - Wheat (<i>Triticum aestivum</i>), - Barley (<i>Hordeum vulgare</i>), - Oats (<i>Avena sativa</i>) - Triticale (<i>x Triticosecale</i>) Wild oats (<i>Avena</i> spp.)			80 to 180 g * # plus an effective rate of a fop herbicide containing fluazifop, haloxyfop, propaquizafop, or quizalofop	# Refer to the label of the partner herbicide for rates and specific directions for use. CropSure Deputy 250WG Herbicide has good activity on barley grass and wild oats but is weaker on brome grass and volunteer cereals so the addition of a partner fop herbicide is generally recommended where any of these other weeds occur together with annual ryegrass. For the partner herbicide use rates at the lowest end of the range recommended for the particular grass weed. Guidance can also be obtained from labelled tank mixes of the partner herbicide with other dim herbicides.

CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Canola (conventional TT, "IMI" varieties)	Annual ryegrass (<i>Lolium rigidum</i>) together with any of the following grasses Annual phalaris (<i>Phalaris spp.</i>), Barley grass (<i>Hordeum spp.</i>), Brome grass (<i>Bromus spp.</i>) Volunteer cereals - Wheat (<i>Triticum aestivum</i>), - Barley (<i>Hordeum vulgare</i>), - Oats (<i>Avena sativa</i>), - Triticale (<i>x Triticosecale</i>), Wild oats (<i>Avena spp.</i>)	2 leaf to early tillering Zadoks 12-23	All States	80 g* # plus an effective rate of a fop herbicide containing fluazifop, haloxyfop, propaquizafop, or quizalofop	Warning: Canola can be sensitive to CropSure Deputy 250WG Herbicide. DO NOT use more than 80 g/ha on canola. To minimise leaf symptoms do not apply to any variety before the majority of plants (80%) are at the true 4 leaf stage (4th leaf expanded 5th leaf emerging). Under certain conditions, all varieties may exhibit leaf symptoms and/or reduced early competitiveness. If this occurs canola yield is usually unaffected but, in some instances, may be reduced. Avoid spraying stressed crops as crops growing under stress will show slower recovery. Avoid spraying practices that lead to over application such as double overlap, and spraying out corners. # Refer to section above for control of additional grasses.
Lucerne Mung beans Navy Beans Peanuts Soybeans	Echinochloa species - Awnless barnyard grass (<i>E. colona</i>), - Barnyard grass (<i>E. crus-galli</i>), - Japanese millet (<i>E. utilis</i>) Digitaria species - Crab grass (<i>D. sanguinalis</i>), - Summer grass (<i>D. ciliaris</i>) Chloris species - Feather top Rhodes grass (<i>C. virgata</i>), - Windmill grass (<i>C. truncata</i>) Setaria species - Dwarf setaria (<i>S. italica</i>), - Whorled pigeon grass (<i>S. verticillata</i>) Brachiaria species - Green summer grass (<i>B. subquadripara</i>), - Velvet grass (<i>B. windersii</i>) Volunteer Crops - Maize (<i>Zea mays</i>), - Sorghum (<i>Sorghum bicolor</i>), Crowsfoot grass (<i>Eleusine indica</i>), Dinebra (<i>Dinebra reflexa</i>), Early spring grass (<i>Eriochloa pseudoacrotricha</i>), Johnson grass (<i>Sorghum halepense</i>) (seedling), Liverseed grass (<i>Urochloa panicoides</i>), Spiny burr grass (<i>Cenchrus incertus</i>)		Qld, NSW, Vic & NT only	120 g or 180 g *	Use the lower rate for the control of seedling grasses at the pre-tillering growth stage and growing under good conditions. Use the higher rate for control of grasses at the early tillering (2 to 3 tillers) growth stage. Aerial application - see Spraying Instructions
Cotton			Qld & NSW only		
Sunflowers			Qld, NSW & Vic only		

CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Sunflowers	Eragrostis species - Elastic grass (<i>E. tenuifolia</i>), - Mexican love grass (<i>E. mexicana</i>), - Stink grass (<i>E. cilianensis</i>)	2 leaf to 5 leaf but prior to tillering	Qld, NSW & Vic only	120g *	For the control of pre-tillering grasses only. Do not apply to tillered grasses
	Volunteer cereals - Wheat (<i>Triticum aestivum</i>), - Barley (<i>Hordeum vulgare</i>)	2 leaf to early tillering Zadoks 12-22, prior to stem elongation		180 g*	For the control of grasses from 2 leaf to early tillering only (max. 2 tillers), prior to stem elongation or booting.
	Other Grasses Coast button grass (<i>Dactyloctenium aegyptium</i>), Grader grass (<i>Themeda Quadrivalvis</i>)	2 leaf to 5 leaf but prior to tillering		180 g*	Use for suppression only of pre-tillered grasses. Do not apply to tillered weeds.

* Always apply with oil spray adjuvant (for example CropSure Takedown Spray Oil) at 1L/100L of spray solution. For aerial application apply oil spray adjuvant at a rate of 1 L/ha.

Note: Does NOT control winter grass (*Poa annua*), Silver grass (*Vulpia* spp.), Nutsedge (*Cyperus* spp.) and broadleaf weeds.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Harvest

NOT REQUIRED WHEN USED AS DIRECTED

Grazing

Canola, Chick peas, Clover, Faba beans, Field peas, Lentils, Linseed, Lucerne, Lupins, Medics, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers, Vetches:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

Cotton:

DO NOT GRAZE OR CUT FOR STOCKFEED. DO NOT FEED COTTON TRASH TO LIVESTOCK.

GENERAL INSTRUCTIONS

Resistant Weeds Warning

GROUP	1	HERBICIDE

CropSure Deputy 250WG Herbicide is a member of the Cyclohexanedione (DIM) group of herbicides. The product has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management, the product is a Group 1 herbicide.

Some naturally occurring weed biotypes resistant to the product and other Group 1 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 1 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Deputy 250WG Herbicide to control resistant weeds.

Stress conditions

Moisture stress is a particular problem, whether caused by drought or short-term stresses, resulting from a combination of high temperatures and low humidity or by prior use of pre-emergence herbicides resulting in stunted root growth of weeds. An irrigation or effective rainfall must occur immediately before or after spraying if these conditions exist. Other stress conditions to avoid include use of pre-emergence herbicides resulting in stunted root growth of weeds, frost, waterlogging and extended cold conditions. Do not spray if any stress conditions are evident just prior to application.

Mixing

Add the required amount of CropSure Deputy 250WG Herbicide (see table) directly to the spray tank which should be 50% full with clean water. After the product is fully dispersed complete filling the tank. The agitation system should be running during mixing and spraying. If agitation ceases and settling occurs, resuspend contents thoroughly before spraying. Ensure that all in-line strainer and nozzle screens in the sprayer are 100 mesh or coarser.

Add any other product next.

An oil adjuvant is essential and should be added last to the spray tank.

Oil adjuvant at 1L/100L of water recommended adjuvant for ground application. For aerial application use oil adjuvant at a rate of 1 L/ha.

Oil adjuvant must be added to at least 10 times its volume of water especially where a suction (filling) probe is used.

The area covered per measure pack will depend upon the application rate/ha according to the following table:

CropSure Deputy 250WG Herbicide Rate/ha	Hectares covered by					
	1 kg	2 kg	3 kg	4 kg	5 kg	10 kg
80 g	12.5	25.0	37.5	50.0	62.5	125
100 g	10.0	20.0	30	40.0	50.0	100
150 g	6.7	13.3	20.0	26.7	33.3	66.7
180 g	5.6	11.1	16.7	22.2	27.8	55.6

Application

Time of application: CropSure Deputy 250WG Herbicide is rain fast within 30 minutes. Do not apply CropSure Deputy 250WG Herbicide if rain is expected within 30 minutes. To obtain the maximum benefits of this product, weeds should be sprayed when young and actively growing. Use the lower rates for grasses with up to 4 leaves. Use the higher rates for weeds with up to 3 tillers.

Unless otherwise specified in the Critical Comments application may be made at any stage of crop growth but allowing sufficient time for the prescribed withholding period.

Method of application: Good spray coverage is essential for maximum results. Spray equipment must be checked and calibrated accurately prior to application.

Boom spraying: Check height of boom above the target weeds to ensure the spray is evenly distributed and a double overlap pattern is obtained. An even distribution of fine droplet size is desirable for maximum efficacy. The inclusion of Supercharge coarsens the spray quality when applied through a TeeJet XR11002 nozzle such that a Medium spray quality results which will help reduce the likelihood of drift.

Broadacre use: Water volumes per hectare will depend on nozzle selection and ground speed but should be in the range of 50-100L/ha. 110° flat fan nozzles are preferred for use with this product. Use higher volumes of spray to achieve better coverage of dense, vigorous weed infestations and/or if the nozzles used produce coarser than a Fine spray quality (measured with water). The use of air induction nozzles is not recommended until further trial data is available.

Aerial application: The product may be applied through boom or Micronair units in 20-30L of water per hectare. The use of the higher volume is preferred due to more reliable results. Spray quality should be Fine droplet size category and swath width should not be wider than recommended for type of plane and application unit. Weed control following aerial application may not be as good as from ground application.

Sprayer clean-up or decontamination

Before spraying sensitive crops (which include canola, chickpeas, faba beans, field peas, lupins, lucerne, clover etc.) thoroughly remove all traces of SU herbicides such as metsulfuron methyl, chlorsulfuron or triasulfuron or Carfentrazone from mixing and spray equipment immediately after use as recommended on the label for that herbicide.

Compatibility

The following recommendations are based on results from field and laboratory tests with CropSure Deputy 250WG Herbicide plus 1% Supercharge. Compatibility is defined as acceptable physical mixing, nil or generally transient crop effect and grass control within 10% of that from CropSure Deputy 250WG Herbicide alone. Performance of any tank mix will be better if the maximum rate of CropSure Deputy 250WG Herbicide is used and the weed growth stages are according to the label. Always refer to the label of the partner product for rates and any specific guidelines and restraints for the crop or situation.

CropSure Deputy 250WG Herbicide is compatible with any one of the following herbicides for the crops specified:

Canola - atrazine, simazine, triasulfuron, herbicides containing fluazifop, haloxyfop, propaquizafop, quizalofop

Pasture, clover based (not brome grass) -

flumetsulam, bromoxynil, MCPA LV (max rate 0.5 L/ha; some clover vigour reduction possible), diflufenican, diflufenican/bromoxynil, simazine, herbicides containing fluazifop, haloxyfop, propaquizafop, quizalofop

CropSure Deputy 250WG Herbicide is compatible with any one of the following insecticides for the crops specified:

Pulses, canola, pasture - alpha-cypermethrin, phosmet, lambda-cyhalothrin, pirimicarb, dimethoate, methidathion

CropSure Deputy 250WG Herbicide is compatible with any one of the following fungicides for the crops specified:

Pulses - procymidone

CropSure Deputy 250WG Herbicide is compatible with any one of the following trace elements for the crops specified:

Canola, Pulses, Pasture - EDTA chelates of Cu, Mn, Zn; sulphates of Cu, Mn, Zn (good quality, such as Agrichem Balance range)

Do not tank mix CropSure Deputy 250WG Herbicide plus oil spray adjuvant with more than one of the above products before doing a jar test beforehand. However, physical compatibility does not guarantee biological compatibility. Do not tank mix with other products or trace elements without reference to an CropSure Pty Ltd representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT plant cereal crops for a period of 4 weeks after application of CropSure Deputy 250WG Herbicide.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to algae and aquatic plants. DO NOT contaminate dams, rivers or streams with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store in prolonged periods of direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling dust. When preparing product for use wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126; New Zealand 0800 764 766.

SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet, which is available from the supplier

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.