

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure **Clethodim 360EC** **Herbicide**

ACTIVE CONSTITUENT: 360 g/L CLETHODIM
SOLVENT: 534 g/L LIQUID HYDROCARBONS

GROUP	1	HERBICIDE
-------	---	-----------

For the control of certain grass weeds in beetroot, cabbage, canola, celery, cotton, forestry, lettuce, non-bearing fruit trees, onions, ornamentals, peanuts, pulses (including azuki beans, broad beans, chickpeas, faba beans, field peas, lentils, lupins and mung beans), potatoes, soybeans and pasture legume (lucerne, clover and medic), seed crops (and pastures) as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT

CONTENTS: 1 L – 1000 L

CropSure Pty Ltd
18 Raymond Road
Laverton North
VIC 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well ventilated area out of direct sunlight. Protect from frost.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, State or Territory government regulations. DO NOT burn empty container or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length chemical resistant gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN A TRANSPORT EMERGENCY
DIAL 000

POLICE OR FIRE BRIGADE

24HR Emergency Response:
AU +61 1800 951 288
NZ +64 800 700 112

APVMA Approval Number: 91580/132710

Batch No:

DOM:

Combustible liquid. Repeated exposure may cause skin dryness or cracking. Harmful if swallowed. May be fatal if swallowed and enters airways. May cause an allergic skin reaction. Harmful to aquatic life with long lasting effects. Keep away from heat, sparks, open flames and hot surfaces. - No smoking. Avoid breathing fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves, protective clothing and eye or face protection. Wash contaminated clothing before reuse. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN: Wash with plenty of soap and water. IF ON SKIN (or hair): Remove immediately all contaminated clothing. Rinse skin with water. If skin irritation or rash occurs: Get medical advice. In case of fire: Use carbon dioxide, dry chemical, foam, to extinguish. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure **Clethodim 360EC** **Herbicide**

ACTIVE CONSTITUENT: 360 g/L CLETHODIM
SOLVENT: 534 g/L LIQUID HYDROCARBONS

GROUP 1 HERBICIDE

For the control of certain grass weeds in beetroot, cabbage, canola, celery, cotton, forestry, lettuce, non-bearing fruit trees, onions, ornamentals, peanuts, pulses (including azuki beans, broad beans, chickpeas, faba beans, field peas, lentils, lupins and mung beans), potatoes, soybeans and pasture legume (lucerne, clover and medic), seed crops (and pastures) as per the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

APVMA APPROVAL NO: 91580/132710

CropSure Pty Ltd
18 Raymond Road
Laverton North
VIC 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

DIRECTIONS FOR USE**RESTRAINTS:**

DO NOT apply without the addition of an adjuvant.

DO NOT apply to plants that are stressed by moisture or temperature extremes.

DO NOT apply if rain is expected within one hour of application.

DO NOT apply CropSure Clethodim 360EC Herbicide more than once to any one crop.

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	RATE mL/ha	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Canola Pulse crops (including azuki beans, broad beans, chickpeas, faba beans, field peas, lentils*, lupins#)	Canola: DO NOT APPLY after flower buds become visible (green buds)	Annual ryegrass (<i>Lolium rigidum</i>), Annual phalaris (<i>Phalaris minor</i>)	100 to 330	2-leaf to fully tillered	Qld, NSW, ACT, Vic, Tas, SA, WA only	Always apply with an adjuvant (for example CropSure Motivate Spray Adjuvant) at 1L/100L or Kwickin* at 1L/100L or a paraffinic spray oil (for example CropSure Takedown Spray Oil) at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides.
	Azuki beans: DO NOT APPLY after first flower buds are visible.	Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus diandrus</i>), Wild Oats (<i>Avena spp.</i>)	115 to 330			
	Chickpeas: DO NOT APPLY beyond full flowering	Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	130 to 330 (115 to 330 in WA only)			
	Faba beans, broad beans: DO NOT APPLY beyond full flowering	Volunteer barley (<i>Hordeum vulgare</i>)	330			
	Field peas: DO NOT APPLY beyond full flowering	Silver grass (<i>Vulpia bromoides</i>) – suppression only	165 to 330		NSW, ACT, Vic, Tas only	
	Lentils: Apply up to the 7 node/early branching stage of crop growth	Paradoxa grass (<i>Phalaris paradoxa</i>)	165 to 250	2- to 5-leaf stage	Qld, NSW, ACT, only "Application up to 7-node/early branching crop growth stage only. # NOT QLD	
	Lupins: DO NOT APPLY after 80% of flowers have opened		250 to 330	2-leaf to fully tillered		
Cotton (Qld, NSW and NT only),	Cotton: DO NOT APPLY after full	Barnyard grass (<i>Echinochloa spp.</i>)	165 to 250	2- to 5-leaf stage	Qld, NSW, ACT, Vic,	Always apply with an adjuvant (for example CropSure Motivate Spray Adjuvant) at
			250 to 330	5-leaf to fully tillered		

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	RATE mL/ha	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Peanuts, Mung beans, Soybeans Pasture legume (lucerne, clover and medic) Seed Crops and Pastures¹ (including pastures containing chicory and plantain).	flowering (mid bloom) Peanuts: DO NOT APPLY after the pod fill stage of crop development Mung beans: DO NOT APPLY after first flower buds are visible Soybeans: DO NOT APPLY after first flower buds are visible	Blown grass (<i>Agrostis aveacea</i>), Crowsfoot grass (<i>Eleusine indica</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass ² (<i>Sorghum halepense</i>), Summer grass (<i>Digitaria</i> spp.), Volunteer sorghum (<i>Sorghum</i> spp.)			WA, NT only	1L/100L or Kwickin* at 1L/100L or a paraffinic spray oil (for example CropSure Takedown Spray Oil) at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides. ¹ See use of CropSure Clethodim 360EC Herbicide in pastures under General Instructions. ² Only Johnson grass seedlings germinating from seed will be controlled. Seedlings that appear later as shoots from underground stems will not be controlled, and more than one application may be required to control emerging shoots.

CROP	WEEDS CONTROLLED	RATE mL/ha	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Beetroot, Cabbage, Celery, Lettuce, Potatoes, Onions	Barnyard grass (<i>Echinochloa</i> spp.), Blown grass (<i>Agrostis aveacea</i>), Crowsfoot grass (<i>Eleusine indica</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Paradoxa grass (<i>Phalaris paradoxa</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass (<i>Sorghum halepense</i>) ¹ , Summer grass (<i>Digitaria</i> spp.), Volunteer sorghum (<i>Sorghum</i> spp.)	165 to 250 250 to 330	2- to 5-leaf stage 5-leaf to fully tillered	All States	Always apply with an adjuvant (for example CropSure Motivate Spray Adjuvant) at 1L/100L or Kwickin* at 1L/100L or a paraffinic spray oil (for example CropSure Takedown Spray Oil) at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. Use a spray volume of 150L/ha when spraying dense grass populations. ¹ Only Johnson grass seedlings germinating from seed will be controlled. Seedlings that appear later as shoots from underground stems will not be controlled, and more than one application may be required to control emerging shoots.
	Annual ryegrass (<i>Lolium rigidum</i>), Annual phalaris (<i>Phalaris minor</i>)	100 to 330	2-leaf to fully tillered	All States	Always apply with an adjuvant (for example CropSure Motivate Spray Adjuvant) at 1L/100L or Kwickin* at 1L/100L or a paraffinic spray oil (for example CropSure Takedown Spray Oil) at 500mL/100L spray volume.
	Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus diandrus</i>), Wild oats (<i>Avena</i> spp.)	115 to 330			The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. Use a spray volume of 150 L/ha when spraying dense grass populations.
	Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	130 to 330 (115 to 330 in WA only)			
	Volunteer barley (<i>Hordeum vulgare</i>)	330			
	Silver grass (<i>Vulpia bromoides</i>) – suppression only (not Qld, WA)	165 to 330			
	Winter grass (<i>Poa annua</i>)	330			

CROP	WEEDS CONTROLLED	RATE mL/ha	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Forestry, Non-bearing Fruit Trees and Ornamentals	Annual ryegrass (<i>Lolium rigidum</i>), Annual phalaris (<i>Phalaris minor</i>), Barley grass (<i>Hordeum leporinum</i>), Barnyard grass (<i>Echinochloa</i> spp.), Blown grass (<i>Agrostis aveacea</i>), Brome grass (<i>Bromus diandrus</i>), Crowsfoot grass (<i>Eleusine indica</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Paradoxa grass (<i>Phalaris paradoxa</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass (<i>Sorghum halepense</i>) ¹ , Silver grass (<i>Vulpia bromoides</i>), – suppression only (not QLD, WA), Summer grass (<i>Digitaria</i> spp.), Volunteer sorghum (<i>Sorghum</i> spp.), Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>), Volunteer barley (<i>Hordeum vulgare</i>), Winter grass (<i>Poa annua</i>)	330	2-leaf to fully tillered	All States	<p>Apply after plants have recovered from transplant shock and are showing signs of active growth.</p> <p>Always apply with an adjuvant (for example CropSure Motivate Spray Adjuvant) at 1L/100L or Kwickin* at 1L/100L or a paraffinic spray oil (for example CropSure Takedown Spray Oil) at 500mL/100L spray volume.</p> <p>Use a spray volume of 150 L/ha when spraying dense grass populations.</p> <p>Do NOT use on Gymnosperms (pines, conifers, etc) unless a prior test has been conducted to check safety on the relevant species.</p> <p>See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for particular species.</p> <p>¹ Only Johnson grass seedlings germinating from seed will be controlled. Seedlings that appear later as shoots from underground stems will not be controlled, and more than one application may be required to control emerging shoots.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST:

PASTURE LEGUME (LUCERNE, CLOVER AND MEDIC) SEED CROPS AND PASTURES: NOT REQUIRED WHEN USED AS DIRECTED.

CANOLA, COTTON, PEANUTS, PULSES (AZUKI BEANS, BROAD BEANS, CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS, MUNG BEANS) AND SOYBEANS: NOT REQUIRED WHEN USED AS DIRECTED.

FORESTRY: NOT REQUIRED WHEN USED AS DIRECTED.

BEETROOT, CABBAGE: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.

ONIONS: DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST.

LETTUCE, POTATOES: DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST.

CELERY: DO NOT APPLY LATER THAN 9 WEEKS BEFORE HARVEST

GRAZING:

PASTURE LEGUME (LUCERNE, CLOVER AND MEDIC) SEED CROPS AND PASTURES: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.

CANOLA, PEANUTS, PULSES (AZUKI BEANS, BROAD BEANS, CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS, MUNG BEANS) AND SOYBEANS: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER APPLICATION.

COTTON: DO NOT GRAZE OR CUT COTTON FORAGE OR STUBBLE FOR STOCK FEED.

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING:

GROUP	1	HERBICIDE
CropSure Clethodim 360EC Herbicide	1	CropSure Clethodim 360EC Herbicide

CropSure Clethodim 360EC Herbicide is a member of the cyclohexanedione group of herbicides. CropSure Clethodim 360EC Herbicide has the inhibition of acetyl coA carboxylase mode of action. For weed resistance management, CropSure Clethodim 360EC Herbicide is a Group 1 Herbicide. Some naturally-occurring weed biotypes resistant to CropSure Clethodim 360EC Herbicide and other Group 1 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CropSure Clethodim 360EC Herbicide or other Group 1 Herbicides. Since occurrence of resistant individuals is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Clethodim 360EC Herbicide to control resistant weeds.

CLEANING SPRAY EQUIPMENT

Before using CropSure Clethodim 360EC Herbicide

Ensure that the recommended clean-out procedure for the previous product (particularly sulfonylurea herbicides) sprayed with the equipment was done properly.

After using CropSure Clethodim 360EC Herbicide

Empty the tank and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean all filters in the tank, pump line and nozzles. Use of a household detergent will aid in cleaning the equipment. Add detergent to the part-filled spray tank and thoroughly circulate through pumps, hoses and nozzles. Drain the system and thoroughly rinse twice with clean water.

Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

MIXING:

To ensure even mixing, half-fill the spray tank with clean water and add the required amount of product. Add spray additive and agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

APPLICATION:

Ground Application:

CropSure Clethodim 360EC Herbicide should be applied with calibrated spray equipment producing MEDIUM spray droplets. Apply in a minimum of 50 litres of water per hectare. Use 150 L/ha when spraying dense populations.

Aerial Application:

CropSure Clethodim 360EC Herbicide can be applied through aircraft fitted with boom or Micronair equipment. A spray volume of 20 to 30 L/ha is recommended and equipment should be adjusted to produce MEDIUM spray droplets.

Best results will be obtained when aerial applications are made in a light crosswind. Applications should not be made during temperature inversions or in conditions of very low relative humidity. Care should be taken to avoid drift damage to adjoining grass crops.

USE OF CROPSURE CLETHODIM 360EC HERBICIDE ON PASTURE LEGUME (LUCERNE, CLOVER AND MEDIC) SEED CROPS AND PASTURES AND PASTURE HERBS:

CropSure Clethodim 360EC Herbicide has been demonstrated to be safe when applied to the following cultivars during early establishment.

Pasture Legumes:**Clover (*Trifolium spp.*)**

Annual: Subterranean clover (*Dalkeith*, *Dalsa*, *Goulburn*, *Koala*, *Nungarrin*, *Riverina*, *Seaton Park and York A*), Arrowleaf clover (*Zulu II*), Balansa clover (*Enduromax*, *Frontier*, *Paradana*), Berseem clover (*Elite II*), Gland clover (*Prima*), Persian clover (*Prolific*, *Leeton*), Rose clover (*Hylon*)
Perennial: Strawberry clover (*Palestine*), White clover (*NuSiral*)

Lucerne and Medics (*Medicago spp.*)

Lucerne (*Aurora*, *Field*, *Genesis*, *Hunterfield*, *Multileaf*, *Salado*, *Sardi SEVEN*, *Sceptre and Venus*), Barrel medic (*Sephi*, *Jester Paraggio*, *Caliph*), Burr medic (*Scimitar*, *Santiago*), Snail medic (*Silver*)

Serradella (*Ornithopus spp.*)

French or Pink serradella (*Cadiz*), Yellow serradella (*Charano*, *Santorini*)

Pasture Herbs:**Chicory (*Puna II*) and Plantain (*Tonic*)**

Apply CropSure Clethodim 360EC Herbicide when plants have reached 2–4 trifoliate leaf stage, or after 40 days from sowing. Some pasture varieties may exhibit an initial reduction in crop vigour or biomass, however, these symptoms are transient and crops recover fully with no effect on yield.

COMPATIBILITY:

CropSure Clethodim 360EC Herbicide is compatible with the following grass herbicides: Targa*, Fusilade*, Haloxyfop (for example CropSure Hallotop 520EC Herbicide), Sertin* and Correct*. (Refer to the Bayer Correct label for detail on adjuvant recommendations for that product).

NOTE: In all situations where CropSure Clethodim 360EC Herbicide is mixed with another grass herbicide, CropSure Clethodim 360EC Herbicide should be used at the full label rate specified for the target weed species. CropSure Clethodim 360EC Herbicide may also be applied in tank mixtures with one of the following products: Sumi-Alpha1 50 EC, Sumi-Alpha1 Dual, Fastac* 100 and Le-Mat*.

CropSure Clethodim 360EC Herbicide should not be applied in a tank mix with Brodal*, Basagran* or Blazer*. Apply as separate applications. For information on compatibility of products not listed, please refer to CropSure Pty Ltd.

CAUTION:

Ornamentals and Trees – While CropSure Clethodim 360EC Herbicide is generally selective to broadleaved plants (i.e., it is active against grasses), tests should always be made on a small number of plants not previously tested. DO NOT use on Gymnosperms (pines, conifers, etc) unless a prior test has been conducted to check safety on the relevant species.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:

DO NOT apply CropSure Clethodim 360EC Herbicide if wind is likely to cause drift onto susceptible crops/plants, cropping lands or pastures. CropSure Clethodim 360EC Herbicide should not be applied through misting equipment or any other method likely to cause excessive drift. Care should be taken to avoid damage to adjoining native grasses or grass crops.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well ventilated area out of direct sunlight. Protect from frost.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, State or Territory government regulations. DO NOT burn empty container or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length chemical resistant gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at ~~his~~ their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.