

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure Beast 200 Herbicide

ACTIVE CONSTITUENT: 200 g/L GLUFOSINATE-AMMONIUM

GROUP	10	HERBICIDE
--------------	-----------	------------------

**For the Non-Residual control of Broadleaf and Grass
Weeds in Various Situations as indicated in the Directions
for Use**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING
OR USING THIS PRODUCT**

Contents: 1L – 1000L

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. May irritate the eyes and repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes. Do not inhale spray mist. When opening the container and preparing spray, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, a washable hat and elbow-length chemical resistant gloves. If applying by boom-spray equipment wear cotton overalls, over normal clothing, buttoned to the neck and wrist elbow-length chemical resistant gloves. In addition, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, a washable hat and half facepiece respirator if applying by low pressure hand wand. Wash hands after use. After each day's use, wash gloves, contaminated clothing and respirator, if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26. New Zealand 0800 764 766.

SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

24HR Medical Emergency Information:

Australia +61 1800 951 288

New Zealand +64 800 700 112

APVMA Approval No.: 90143/127764

Batch No.:

Date of Manufacture:

May cause drowsiness or dizziness. May damage fertility. May damage the unborn child. May cause damage to organs through prolonged or repeated exposure. Harmful if swallowed, in contact with skin or if inhaled. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. Wear protective gloves, protective clothing and eye or face protection. Call a POISON CENTRE or doctor if you feel unwell. Get medical advice or attention if you feel unwell. Wash contaminated clothing before reuse. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove victim to fresh air and keep comfortable for breathing. If exposed or concerned: Get medical advice. In case of fire: Use carbon dioxide, dry chemical, foam, to extinguish. Store locked up. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure Beast 200 Herbicide

ACTIVE CONSTITUENT: 200 g/L GLUFOSINATE-AMMONIUM

GROUP	10	HERBICIDE
--------------	-----------	------------------

**For the Non-Residual Control of Broadleaf and Grass
Weeds in Various Situations as indicated in the Directions
for Use**

**IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING
THIS PRODUCT**

APVMA Approval No.: 90143/127764

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply by aircraft.

DO NOT apply when rain is expected within 6 hours.

DO NOT apply to weeds under stress due to, for example, very dry, very wet, frosty or diseased conditions.

DO NOT apply under hot dry conditions (temperatures above 33°C with a relative humidity below 50%).

Crop / Situation	Weed	State	Rate	WHP	Critical Comments
Blackberry, boysenberry, loganberry, raspberry	Primocane and sucker control	NSW, Vic, Tas only	500mL/ 100L water	Nil	Apply as a directed spray to suckers and primocanes. Contact with flowers, developing fruit or desirable foliage will cause damage. Ensure complete coverage of primocanes/suckers by spraying to the point of runoff, preferably when they are less than 15cm high. Wetting agent may be added at a rate of 25mL/100L or equivalent.
Avocado, banana, feijoa, guava, kiwifruit, litchi, mango, pawpaw, passionfruit, pineapple, rambutan plantations	See list of weeds controlled in Tables 1 and 2.	Qld, NSW, Vic, SA, WA, NT only	1.0 to 5.0 L/ha	Nil	<p>Apply as a directed or shielded spray. Refer to the label section Application Equipment for specific information on application methods. Controlled Droplet Application equipment must not be used for application in cherry orchards.</p> <p>Warnings: DO NOT allow spray or spray drift to contact desirable foliage or green (uncalloused) bark. To avoid potential crop damage, refer to the label sections on Application Equipment and PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.</p> <p>CropSure Beast 200 Herbicide may be used around trees/vines less than two years old provided they are effectively shielded from spray and spray drift.</p> <p>The recommended rate of use is determined by the following criteria:</p> <ul style="list-style-type: none"> WEED SPECIES WEED STAGE OF GROWTH WEED DENSITY CLIMATIC CONDITIONS <p>WEED SPECIES Apply the appropriate rate to control the least susceptible weed present as per the lists of weeds controlled in the accompanying tables.</p> <p>WEED STAGE OF GROWTH Use the lower rate when weeds are young and succulent (grasses: pre-tillering; broadleaves: cotyledons to 4-leaf) or the population is very sparse. A median rate should be used for medium sized plants (grasses: tillering; broadleaves: 4-leaf to advanced vegetative) and the high rate should be used when weeds are mature (grasses: nodding to flowering; broadleaves: budding to flowering).</p>
Citrus orchards		All States			
Olive plantations					
Pome and stone fruit orchards			21 days (H)		
Tree nut plantations Vineyards				Nil	

Crop / Situation	Weed	State	Rate	WHP	Critical Comments
				-	<p>WEED DENSITY Use the higher rates when the weed population is dense. Thorough coverage of weeds is essential for good control.</p> <p>CLIMATIC CONDITIONS Best results are achieved when applied under warm humid conditions. Control will be reduced and/or slower under cold conditions and/or overcast conditions. Good results will be achieved under most other conditions, however poor results may occur under hot dry conditions (temperature above 33°C with a relative humidity below 50%). Weeds that have been hardened or stunted in growth due to stressed conditions should be treated at the maximum rate.</p> <p>COVERAGE Complete coverage of weeds is essential for good control. Poor coverage may result in re-growth.</p> <p>PERENNIAL WEEDS Apply when weeds are actively growing. Follow-up treatments will be necessary to control re-growth of perennial weeds in most cases.</p>
Strawberries, cane berry fruits (inter-row)	See lists of weeds controlled in Tables 1 and 2	All states	1.0 to 5.0 L/ha	Nil	Apply as a directed or shielded spray to the inter-row area. Take care not to allow spray or spray drift to contact the crop, including strawberry runners. Refer to GENERAL INSTRUCTIONS for warnings concerning plastic mulch and fumigated/sterilised soil. Determine the recommended rate of use by considering the criteria WEED SPECIES, WEED STAGE OF GROWTH, WEED DENSITY and CLIMATIC CONDITIONS, as described above.
Tomatoes (inter-row)					
Commercial & Industrial areas, rights-of-way and other non-agricultural areas		All states	1.0 to 6.0 L/ha	-	Determine the recommended rate of use by considering the criteria WEED SPECIES, WEED STAGE OF GROWTH, WEED DENSITY and CLIMATIC CONDITIONS as described above. Warnings: Do not allow spray or spray drift to contact desirable plants. To avoid potential crop damage, refer to the label sections on Application Equipment and PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIOD (WHP)

HARVEST (H)

avocado, banana, feijoa, guava, kiwifruit, litchi, mango, olives, pawpaw, passionfruit, pineapple, rambutan, blackberry, boysenberry, loganberry, raspberry, citrus fruit, grapes, strawberries, tomatoes, tree nuts: **NOT REQUIRED WHEN USED AS DIRECTED**

Pome and stone fruit – **DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION**

GRAZING (G)

DO NOT GRAZE OR CUT TREATED AREAS FOR STOCKFOOD FOR 8 WEEKS AFTER APPLICATION

Table 1. Recommendations for weed control (except when referred to Table 2).

Common Name	Scientific Name	Application Rates		
		Boom or Directed Sprayer L/ha	Handgun mL/100L	Knapsack mL/15L
ANNUAL WEEDS				
Amaranthus spp.	<i>Amaranthus spp.</i>	2.0 to 5.0	500	75
Apple of Peru	<i>Nicandra physalodes</i>	1.5 to 3.0	300	45
Argentine peppercress	<i>Lepidium bonariense</i>	2.0 to 3.0	300	45
Awnless barnyard grass	<i>Echinochloa colona</i>	2.5 to 3.5	350	53
Barley grass	<i>Hordeum leporinum</i>	2.0 to 3.0	300	45
Barnyard grass	<i>Echinochloa crus-galli</i>	2.0 to 5.0	500	75
Billy goat weed	<i>Ageratum conyzoides</i>	2.0 to 5.0	500	75
Bitter cress	<i>Cardamine hirsute</i>	2.0 to 5.0	500	75
Black bindweed (buckwheat) (refer Note 2)	<i>Fallopia convolvulus</i>	1.8 to 5.0	500	75
Bladder ketmia	<i>Hibiscus trionum</i>	3.0 to 5.0	500	75
Bordered panic	<i>Entolasia marginata</i>	2.0 to 4.0	400	60
Brome grass (refer Note 1)	<i>Bromus spp.</i>	2.0 to 3.0	300	45
Calopo	<i>Calopogonium mucanoides</i>	2.0 to 5.0	500	75
Caltrop burr (refer also Table 2)	<i>Tribulus terrestris</i>	3.0 to 5.0	500	75
Capeweed	<i>Arctotheca calendula</i>	1.5 to 5.0	500	75
Clover (subterranean)	<i>Trifolium subterranean</i>	1.8 to 3.0	300	45
Cobbler's peg	<i>Bidens pilosa</i>	2.0 to 5.0	500	75
Common storksbill	<i>Erodium cicutarium</i>	1.5 to 4.0	400	60
Crowsfoot grass	<i>Eleusine indica</i>	3.0 to 5.0	500	75
Deadnettle (refer also Table 2)	<i>Lamium amplexicaule</i>	2.0 to 5.0	500	75
Dwarf crumbweed	<i>Chenopodium pumilio</i>	3.0 to 5.0	500	75
Fat hen	<i>Chenopodium album</i>	3.0 to 5.0	500	75
Fumitory	<i>Fumaria officinalis</i>	1.8 to 5.0	500	75
Green crumbweed	<i>Chenopodium carinatum</i>	2.0 to 5.0	500	75
Lesser canary grass (refer also Table 2)	<i>Phalaris minor</i>	3.0 to 5.0	500	75
Liverseed grass (refer also Table 2)	<i>Urochloa panicoides</i>	1.5 to 5.0	500	75
Medics (annual)	<i>Medicago spp.</i>	1.0 to 5.0	500	75
Milk thistle	<i>Sonchus oleraceus</i>	2.0 to 5.0	500	75
Mint weed	<i>Salvia reflexa</i>	3.0 to 5.0	500	75
New Zealand spinach	<i>Tetragonia tetragonoides</i>	2.0 to 5.0	500	75
Patterson's Curse	<i>Echium plantagineum</i>	1.0 to 3.0	300	45
Peanuts	<i>Arachis hypogaea</i>	1.5 to 3.0	300	45
Pigweed	<i>Portulaca oleracea</i>	3.0 to 5.0	500	75
Pinkbur	<i>Urena lobata</i>	2.0 to 5.0	500	75
Potato weed	<i>Galinsoga parviflora</i>	2.0 to 5.0	500	75
Praire grass (refer Note 1)	<i>Bromus unioloides</i>	4.0 to 5.0	500	75
Prickly lettuce	<i>Lactuca serriola</i>	3.0 to 5.0	500	75
Red natal grass	<i>Rhynchoselytrum repens</i>	2.0 to 5.0	500	75
Ryegrass (annual)	<i>Lolium rigidum</i>	2.0 to 5.0	500	75
Saffron thistle	<i>Carthamus lanatus</i>	1.5 to 5.0	500	75
St. Barnby's thistle	<i>Centaurea solstitialis</i>	1.5 to 5.0	500	75
Sago weed	<i>Plantago cunninghamii</i>	2.0 to 3.0	300	45
Scarlet pimpernel	<i>Anagallis arvensis</i>	2.0 to 5.0	500	75
Setaria	<i>Setaria italica</i>	2.0 to 5.0	500	75
Sheep thistle	<i>Carduus tenuiflorus</i>	2.5 to 5.0	500	75
Silver grass	<i>Vulpia myuros</i>	2.0 to 5.0	500	75
Sorghum/sudax	<i>Sorghum bicolor</i>	2.0 to 5.0	500	75
Square weed	<i>Spermacoce latifolia</i>	2.0 to 5.0	500	75
Stagger weed	<i>Stachys arvensis</i>	2.0 to 5.0	500	75
Star of Bethlehem	<i>Ipomoea quamoclit</i>	2.0 to 5.0	500	75
Summer grass	<i>Digitaria ciliaris</i>	2.0 to 5.0	500	75
Thickhead	<i>Crassocephalum crepidioides</i>	3.0 to 5.0	500	75
Three Cornered Jack	<i>Emex australis</i>	2.0 to 5.0	500	75
Tomato	<i>Lycopersicon esculentum</i>	2.0 to 5.0	500	75
Turnip weed	<i>Rapistrum rugosum</i>	3.0 to 5.0	500	75

Common Name	Scientific Name	Application Rates		
		Boom or Directed Sprayer L/ha	Handgun mL/100L	Knapsack mL/15L
Variegated thistle (refer also Table 2)	<i>Silybum marianum</i>	2.5 to 5.0	500	75
Wheat	<i>Triticum aestivum</i>	4.0 to 5.0	500	75
Wild carrot	<i>Daucus glochidiatus</i>	2.0 to 5.0	500	75
Wild gooseberry	<i>Physalis minima</i>	2.0 to 5.0	500	75
Wild mustard	<i>Sisymbrium orientale</i>	2.0 to 5.0	500	75
Wild oats (refer also Table 2)	<i>Avena spp.</i>	3.0 to 5.0	500	75
Wild radish	<i>Raphanus raphanistrum</i>	5.0	500	75
Wire weed (refer also Table 2)	<i>Polygonum aviculare</i>	1.5 to 5.0	500	75
PERENNIAL WEEDS				
Blady grass	<i>Imperata cylindrica</i>	3.0 to 4.0	400	60
Cape tulip	<i>Homeria spp.</i>	2.0 to 3.0	300	45
Centro	<i>Centrosema pubescens</i>	1.0 to 5.0	500	75
Clover glycine	<i>Glycine latrobeana</i>	1.0 to 3.0	300	45
Couch grass	<i>Cynodon dactylon</i>	2.5 to 5.0	500	75
Cow pea	<i>Vigna unguiculata</i>	1.0 to 3.0	300	45
Giant sensitive plant	<i>Mimosa invisa</i>	2.0 to 5.0	500	75
Greenleaf desmodium	<i>Desmodium intortum</i>	1.0 to 3.0	300	45
Johnson grass	<i>Sorghum halepense</i>	3.0 to 5.0	500	75
Panicum spp.	<i>Panicum spp.</i>	2.0 to 5.0	500	75
Paspalum spp.	<i>Paspalum spp.</i>	3.0 to 5.0	500	75
Perennial bindweed	<i>Convolvulus arvensis</i>	2.0 to 3.0	300	45
Shamrock	<i>Oxalis corymbosa</i>	3.0	300	45
Sida weed (refer also Table 2)	<i>Sida retusa</i>	3.0 to 5.0	500	75
Silver leaf desmodium	<i>Desmodium uncinatum</i>	4.0 to 5.0	500	75
Siratro	<i>Macroptilium atropurpureum</i>	1.0 to 3.0	300	45
Stink grass	<i>Eragrostis ciliaris</i>	3.0 to 5.0	500	75
White clover	<i>Trifolium repens</i>	3.0 to 5.0	500	75
White eye	<i>Richardia brasiliensis</i>	3.0 to 5.0	500	75
Willow herb	<i>Epilobium spp.</i>	4.0 to 5.0	500	75

Notes:

1. Well-established clumps of Prairie grass and Brome grasses may only be suppressed at these rates. Follow-up treatments may be necessary to control re-growth.
2. Good control will be achieved on small and medium sized plants only in non-crop situation.

Table 2. For control of weeds in Commercial and Industrial areas, rights-of-way and other non-agricultural areas (when referred from Table 1).

Common Name	Scientific Name	Application Rate		
		Boom or Directed Sprayer L/ha	Handgun mL/100L	Knapsack mL/15L
ANNUAL WEEDS				
Caltrop burr	<i>Tribulus terrestris</i>	4.0 to 5.0	500	75
Dead nettle	<i>Lamium amplexicaule</i>	6.0	600	90
Lesser canary grass	<i>Phalaris minor</i>	4.0 to 6.0	600	90
Liverseed grass	<i>Urochloa panicoides</i>	1.5	150	23
Variegated thistle	<i>Silybum marianum</i>	6.0	600	90
Wild oats	<i>Avena spp.</i>	5.0 to 6.0	600	90
Wire weed	<i>Polygonum aviculare</i>	2.0 to 5.0	500	75
PERENNIAL WEEDS				
Sida weed	<i>Sida retusa</i>	4.0 to 5.0	500	75

GENERAL INSTRUCTIONS

CropSure Beast 200 Herbicide is a non-volatile herbicide with activity against many annual and perennial broadleaf weeds and grasses.

CropSure Beast 200 Herbicide is absorbed by plant foliage and green stems. It is not significantly translocated as an active herbicide throughout the plant, and therefore will only kill that part of a green plant that is contacted by spray. CropSure Beast 200 Herbicide does not provide residual weed control. Visible symptoms of control appear in 3 to 7 days, but complete desiccation may take 20 to 30 days under cool conditions.

Best results are achieved when application is made under good growing conditions. Application to weeds under stress (e.g. due to continuous severe frosts, dry or waterlogged conditions) should be avoided.

Soil fumigation / sterilisation

CropSure Beast 200 Herbicide is metabolised (broken down) by microorganisms in the soil to become inactive. Soil fumigation or sterilisation will reduce the number of microorganisms present, thus slowing the breakdown of CropSure Beast 200 Herbicide. As damage to transplants or seedlings may occur, it is not advisable to apply CropSure Beast 200 Herbicide in conjunction with soil fumigation or sterilisation.

Plastic mulches

CropSure Beast 200 Herbicide will remain active on inert surfaces such as plastic. Special care should be taken when applying CropSure Beast 200 Herbicide over plastic mulches, as plant contact with the mulch after spraying may result in crop damage.

RESISTANT WEEDS WARNING

GROUP	10	HERBICIDE
-------	----	-----------

CropSure Beast 200 Herbicide is a member of the glycine group of herbicides. CropSure Beast 200 Herbicide has the inhibitor of glutamine synthetase mode of action. For weed resistance management CropSure Beast 200 Herbicide is a Group 10 herbicide.

Some naturally occurring weed biotypes resistant to CropSure Beast 200 Herbicide, and other Group 10 herbicides which inhibit glutamine synthetase, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CropSure Beast 200 Herbicide or other Group 10 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Beast 200 Herbicide to control resistant weeds.

Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with CropSure Beast 200 Herbicide. If you are growing produce for export, please check with CropSure Pty Ltd for the latest information on MRLs and import tolerances BEFORE using CropSure Beast 200 Herbicide.

Compatibility

CropSure Beast 200 Herbicide is compatible with most residual herbicides e.g. simazine, oxyfluorfen, norfluazuron, and oryzalin, and with glyphosate and metsulfuron. The addition of a wetting agent or other adjuvant is generally not considered necessary, (refer to the Directions for Use table). However, benefit has been obtained using a wetting agent or adjuvant on hard-to-wet weeds when using water rates in excess of 500 L/ha. The rate is 25 mL/100 L of a 1000 g/L non-ionic wetting agent, or equivalent. For information on compatible wetting agents and adjuvants, contact your local CropSure Pty Ltd representative.

Mixing

CropSure Beast 200 Herbicide mixes easily with water. Clean water should always be used for mixing with CropSure Beast 200 Herbicide. Ensure that the spray tank is free of any residues of previous spray materials. Two-thirds fill the spray tank with clean water, and with agitator operating add the required amount of CropSure Beast 200 Herbicide. Add other relevant compatible products. Top the tank up to the required volume with clean water with agitator running.

Application Equipment**Ground Sprayers**

Aim to apply a thorough and even coverage of spray to the target plant. Dense stands of weeds should be thoroughly wetted with spray. Incomplete coverage may result in poor control. Equipment should be such that adequate coverage, penetration and volume of spray liquid can be achieved.

Boom or Directed Sprayer Equipment

CropSure Beast 200 Herbicide should be applied at label rates (refer to specific column in the lists of weeds controlled) in sufficient water to give thorough coverage of weeds. It has been found that 300 to 500 L/ha has given good results under most weed conditions. Special care must be taken when using sprayer/slasher combination units not to cause dust and turbulence, which can carry spray into non-target areas.

Knapsack and Handgun Equipment

CropSure Beast 200 Herbicide should be applied at label rates (refer to specific columns in the lists of weeds controlled) in adequate water to thoroughly wet the weeds being sprayed, i.e. 500 to 1000 L/ha. Dense stands will require up to 1000 L/ha of spray mixture, whereas less dense stands will require less water. High volume application using hollow-cone nozzles for hand spraying is recommended.

Controlled Droplet Application (CDA) Equipment

CropSure Beast 200 Herbicide may be applied through CDA row spraying equipment fitted with a solid (impermeable) shroud or skirt, at rates as recommended for boom or directed sprayers (refer to specific column in the lists of weeds controlled), provided thorough spray coverage of weeds can be achieved. Apply preferably when weeds are less than 15 cm in height, with the equipment set up so that the spray dome only just touches the tops of the weeds. A total spray volume of 20 to 30 L/ha has been found to give good results. Do not mix residual herbicides or any spray adjuvants with CropSure Beast 200 Herbicide when using CDA equipment.

Warning: Because the spray solution is highly concentrated particular care must be taken when using CropSure Beast 200 Herbicide through CDA equipment to avoid contact of the spray solution with any part of the crop trunk or canopy. DO NOT apply CropSure Beast 200 Herbicide through equipment fitted with bristle skirts. Particular care should be taken when using CDA equipment around green or uncalloused bark.

Please refer to PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.

CDA equipment must not be used for application in cherry orchards.

Sprayer clean-up

Clean all equipment after use by thoroughly flushing with water.

Aircraft

Do not apply by aircraft.

PRECAUTIONS**Re-entry period**

Do not allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with this product or the used container.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT apply on desirable foliage or allow spray to drift onto the foliage of desirable plants, trees or vines, as damage will occur.

DO NOT allow product to contact green or uncalloused bark (such as on desirable young trees and vines) or cut, cracked, damaged or wounded tissue, where the affected surface is not adequately healed. CropSure Beast 200 Herbicide may be used around desirable trees/vines less than two years old provided they are effectively shielded from spray and spray drift.

DO NOT allow desirable plant foliage to contact any inert surface, such as plastic mulches, which have been treated with CropSure Beast 200 Herbicide.

DO NOT apply CropSure Beast 200 Herbicide to recently fumigated or sterilised soil.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. May irritate the eyes and repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes. Do not inhale spray mist. When opening the container and preparing spray, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, a washable hat and elbow-length chemical resistant gloves. If applying by boom-spray equipment wear cotton overalls, over normal clothing, buttoned to the neck and wrist elbow-length chemical resistant gloves. In addition, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, a washable hat and half facepiece respirator if applying by low pressure hand wand. Wash hands after use. After each day's use, wash gloves, contaminated clothing and respirator, if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26. New Zealand 0800 764 766.

SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.