

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure **Tebustar 430SC** **Fungicide**

ACTIVE CONSTITUENT: 430 g/L TEBUCONAZOLE

GROUP

3

FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT

Contents: 10L - 1000L

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

DrumMUSTER (10 L, 20 L)

This container can be recycled if it is clean, dry, free of visible residues and has the DrumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any DrumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

For refillable containers (110 L, 1000 L)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

24HR Emergency Response:
AU +61 1800 951 288
NZ +64 800 700 112

APVMA Approval No.: 91537/132555

Batch No:

Date of Manufacture:

Harmful if swallowed. Suspected of damaging the unborn child. Very toxic to aquatic life with long lasting effects. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not get in eyes, on skin, or on clothing. Avoid contact during pregnancy or while nursing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. If exposed or concerned: Get medical advice. In case of fire: Use carbon dioxide, dry chemical, foam, water fog, to extinguish. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure Tebustar 430SC Fungicide

ACTIVE CONSTITUENT: 430 g/L TEBUCONAZOLE

GROUP

3

FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

APVMA Approval No.: 91537/132555

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

DIRECTIONS FOR USE

Crop	Disease	Application	Rate mL/ha	WHP	Critical Comments
Bananas (Qld, NSW, WA, NT only)	Leaf spot (Yellow sigatoka) Leaf speckle Black Sigatoka	Ground Application Apply by misting machine or airblast sprayer in a convenient volume of water Aerial application Apply at least 20L of spray mixture per hectare	230 Add 3-6 L water miscible oil per ha	H 1 day	Maintain good de-leaving practices to reduce disease inoculum. Very old leaves and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of CropSure Tebustar 430SC Fungicide. Tropical Areas (e.g., North Qld, NT, Ord River): Apply a regular schedule of protectant sprays. When conditions favour disease, apply a maximum of 2 consecutive sprays at 14 day intervals. Do not apply more than 6 CropSure Tebustar 430SC Fungicide sprays in any 12 month period. Do not apply any CropSure Tebustar 430SC Fungicide sprays in the months of June, July, August and September. Sub-Tropical Areas (e.g., South Qld, NSW): Commence spraying with CropSure Tebustar 430SC Fungicide at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a maximum of 2 consecutive CropSure Tebustar 430SC Fungicide sprays. Do not apply more than 5 CropSure Tebustar 430SC Fungicide sprays in any 12 month period. CropSure Tebustar 430SC Fungicide is approved for use in banana plantations inter-planted with avocados.
Peanuts (South Qld and NSW only)	Early Leaf spot Late leaf spot Rust Net Blotch	Low Disease Ground Application Apply in at least 100L of water per ha Aerial application Apply at least 30L of spray mixture per hectare	175 + Agridex 1L/ha	H 3 weeks G 3 weeks	Regularly check high-risk areas in the crop for disease (e.g., lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found, then either; <ul style="list-style-type: none">• Spray immediately after last disease conducive weather (e.g., rain or heavy dews), or <ul style="list-style-type: none">• Spray before the next conducive weather. Repeat after 14 days if conditions remain favourable to disease development. If not, repeat just before or just after the next disease conducive weather. For resistance strategy, see General Instructions.
Peanuts (North Qld, WA, NT only)	Early Leaf spot Late leaf spot Rust	Low Disease Moderate disease Severe disease	230 + Agridex 1L/ha 290 + Agridex 1L/ha 440 + Agridex 1L/ha		Begin spraying at 3-4 weeks after planting. If band spraying, apply half the appropriate rate as a 45cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14-day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.
	Net blotch	Low Disease Moderate - severe disease	290 + Agridex 1L/ha 440 + Agridex 1L/ha		Apply at 14 day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.

Crop	Disease	Application	Rate mL/ha	WHP	Critical Comments
Green Beans	Rust	Can be applied by aircraft or ground rig	350 + Agridex 1L/ha	H 3 days G 3 weeks	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high.
Peas	Powdery Mildew	Ground Application Apply in at least 50L of water per ha	145		Apply at flowering or at the first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions.
Wheat	Leaf rust Stripe rust Septoria nodorum blotch Yellow leaf spot	Aerial application Apply in at least 10L of spray mixture per hectare	145 or 290	H 5 weeks G 14 days	Use the higher rate when longer disease control is required. Stripe rust: See spray timings under General Instructions. Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil at 1% may improve performance of CropSure Tebustar 430SC Fungicide on wheat, oats and barley.
Oats	Septoria tritici blotch		290		
Barley	Crown rust		145 or 290		Apply at late tillering to early jointing.
Wheat Oats	Scald		145		Apply when 5% of the leaf area is infected. Use higher rate when longer disease control is required.
Wheat Oats	Powdery mildew		145 or 290		
Onions (Tas only)	White root rot	Before sowing apply CropSure Tebustar 430SC Fungicide onto lime super. Ensure good coverage of all lime super particles.	1.45 mL/100 m of row mixed with 145-218 g lime super/100 m of row	-	Apply CropSure Tebustar 430SC Fungicide treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a bandwidth of 2cm. Ensure that the correct rate of CropSure Tebustar 430SC Fungicide is used otherwise some delay in emergence and reduced stands of seedlings may occur.
Pawpaw	Black spot	Ensure thorough coverage of leaves and fruit.	290	H 3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of CropSure Tebustar 430SC Fungicide. Apply CropSure Tebustar 430SC Fungicide at 14 day intervals. Alternate CropSure Tebustar 430SC Fungicide with sprays of a protectant fungicide (for example CropSure Greenshield 750WG Fungicide). Do not apply more than 6 sprays of CropSure Tebustar 430SC Fungicide (or any DMI fungicide) on any block in any 12 month period.
Pyrethrum	<i>Sclerotinia sclerotiorum</i>		350	-	Apply twice, in rotation with other control measures, at 7-10 day intervals. Commence at 1 to 2% flowering. Use under direction of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf rust Stem rust	Apply in at least 100L of water per ha	290	-	Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or re-infection may require a further application 3 – 4 weeks later. Ensure thorough coverage and use higher water volumes in dense or advanced crops.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS**BANANAS, AVOCADOS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.****BEANS, PEAS: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 3 DAYS AFTER APPLICATION.****CEREALS: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 14 DAYS AFTER APPLICATION.****ONIONS: NOT REQUIRED WHEN USED AS DIRECTED.****PAWPAW: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.****PEANUTS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 21 DAYS AFTER APPLICATION.****H = Harvest, G = Grazing****GENERAL INSTRUCTIONS****FUNGICIDE RESISTANCE WARNING**

GROUP	3	FUNGICIDE
-------	---	-----------

CropSure Tebustar 430SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 Fungicide.

Some naturally occurring individual fungi resistant to CropSure Tebustar 430SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by CropSure Tebustar 430SC Fungicide or other Group 3 Fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Tebustar 430SC Fungicide to control resistant fungi.

RESISTANCE MANAGEMENT RECOMMENDATIONS – PEANUTS

Apply no more than 3 consecutive sprays of DMI fungicides (e.g., tebuconazole) before switching to a non-DMI fungicide. Apply no more than 5 DMI sprays per season.

SPECIAL WARNING – BANANAS

There are certain conditions where the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise risk of fruit marking.

Never include adjuvants other than water miscible oils with CropSure Tebustar 430SC Fungicide sprays. The inclusion of wetting agents is known to cause phototoxicity to young fruit.

FOLIAR DISEASES ON CEREAL CROPS

Do not apply to cereal crops more than once in a season. Treatment will give approximately three weeks disease suppression. Economic response may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7-10 days after application.

Yield potential: crops with potential yield under 2 t/ha are unlikely to give economic responses to a fungicide spray except under conditions of very severe disease. Economic responses are most likely with crops with potential yield of over 3 t/ha.

Spray Timings for Stripe Rust Control

Obtain advisory literature from Department of Agriculture for classification of resistant and susceptible varieties.

The following spray program is suggested:

Seedling infections: When approximately 20 out of 100 leaves show signs of infection during tillering or jointing, apply a spray within a week.

Adult infections

Susceptible varieties: When approximately 10 out of 100 leaves show first sign of infection apply a spray within a week.

Moderately Susceptible varieties: When approximately 15-20 out of 100 leaves show first sign of infection apply a spray within a week. Do not delay.

Moderately Resistant and Resistant Varieties: Monitor carefully. If rust appears and spreads, spray within 1 week.

MIXING

Prior to pouring, shake container vigorously, then add the required amount of CropSure Tebustar 430SC Fungicide to water in the spray vat while stirring or with agitators in motion. Add the required amount of Agridex (peanuts, beans) or water miscible oil (bananas) and mix thoroughly.

APPLICATION

Aircraft should fly as low as possible under the prevailing conditions to minimise drift.

PRECAUTIONS

Re-Entry Period

Do not allow entry into treated areas until the spray has dried.

When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with chemical or used containers. A spray drift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practise Manual.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

DrumMUSTER (10 L, 20 L)

This container can be recycled if it is clean, dry, free of visible residues and has the DrumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any DrumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

For refillable containers (110 L, 1000 L)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.