

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure **Amigo 625** **Herbicide**

ACTIVE CONSTITUENT:
625 g/L 2,4-D present as the dimethylamine and diethanolamine salts

GROUP 4 HERBICIDE

For the control of broadleaf weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, hardwood and soft wood plantations, lucerne, oil tea tree, pastures, sugar cane, peanuts and non agricultural areas as per the directions for use.

THIS IS A PHENOXY HERBICIDE THAT CAN CAUSE SEVERE DAMAGE TO NATIVE VEGETATION AND SUSCEPTIBLE CROPS SUCH AS COTTON, GRAPES, TOMATOES, OILSEED CROPS AND ORNAMENTALS.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT

CONTENTS : 5L - 1000L

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility.

If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

For Mini Bulk Returnable Containers

Store the original sealed container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the MicroMatic valve or the security seal. DO NOT contaminate the container with water or any other foreign matter. After each use of the product ensure the MicroMatic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the container have been used, please return the empty container to the point of purchase. The container remains the property of CropSure Pty Ltd.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact, inhaled or swallowed. Corrosive to the eyes and skin. Will irritate the nose and throat. Avoid contact with the eyes and skin. Do not inhale vapour or spray mist.

When opening the container and preparing spray or using undiluted concentrate, wear chemical resistant waterproof clothing over a layer of normal clothing and a washable hat, elbow-length chemical resistant gloves, impervious footwear and full face piece respirator with organic vapour/gas cartridge or canister.

If applying by boom spray equipment with enclosed operator's cab and air filtration or aerial spraying equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.

If applying by boom spray equipment with open operator's cab or hand-held spray equipment wear chemical resistant waterproof clothing over a layer of normal clothing and a washable hat, elbow-length chemical resistant gloves and full face piece respirator with organic vapour/gas cartridge or canister.

If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves (and) respirator and if rubber wash with detergent and warm water contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766.

SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE

24HR Emergency Response:
AU +61 1800 951 288
NZ +64 800 700 112

APVMA APPROVAL NO: 90999 / 138528

BATCH NO:

DATE OF MANUFACTURE:

Harmful if swallowed. May cause an allergic skin reaction. Causes serious eye damage. Toxic to aquatic life with long lasting effects. Avoid breathing fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves, protective clothing and eye or face protection. Immediately call a POISON CENTRE or doctor/physician. Wash contaminated clothing before reuse. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN: Wash with plenty of soap and water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin irritation or rash occurs: Get medical advice. In case of fire: Use carbon dioxide, dry chemical, foam, water fog, to extinguish. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure **Amigo 625** **Herbicide**

ACTIVE CONSTITUENT:
625 g/L 2,4-D present as the dimethylamine and diethanolamine salts

GROUP 4 HERBICIDE

For the control of broadleaf weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, hardwood and softwood plantations, lucerne, oil tea tree, pastures, sugar cane, peanuts and non agricultural areas as per the directions for use.

THIS IS A PHENOXY HERBICIDE THAT CAN CAUSE SEVERE DAMAGE TO NATIVE VEGETATION AND SUSCEPTIBLE CROPS SUCH AS COTTON, GRAPES, TOMATOES, OILSEED CROPS AND ORNAMENTALS.

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

APVMA Approval No.: 90999 / 138528

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

DIRECTIONS FOR USE**RESTRAINTS:****GENERAL RESTRAINTS**

DO NOT exceed maximum application rate of 7.2L/ha (4500 g ae/ha).

DO NOT apply by hand-held spraying equipment at rates exceeding 5.3 L/ha or 530 mL/100 L.

DO NOT apply by spraying equipment carried on the back of the user (Manually pressurised backpack sprayer).

DO NOT apply by aircraft at rates exceeding 3.6 L/ha.

DO NOT use open mixing/loading equipment if treating more than 50 hectares in one day.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of runoff for at least 3 days after application.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

Additional USAGE restrictions apply in some crops, states and seasons, see restriction tables 1, 2, 3, 4 and 5.

Table 1: Timing restrictions for spraying peanuts

Situation	Rate (L/ha)	Region	Timing Restriction
DO NOT APPLY DURING THE MONTHS			
Broadcast spraying, prior to sowing (peanuts)	Up to 1.4L/ha	Cape York	October and November
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	No timing restrictions
		Burdekin	October
		Mackay/Whitsunday	September to December
		Mary/Burnett	October to November
		SE Queensland	August to May
	Up to 1.7 L/ha	Cape York	October and November
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	No timing restrictions
		Burdekin	October
		Mackay/Whitsunday	August to December
		Mary/Burnett	September to November
		SE Queensland	Use not supported
Band spraying, post-sowing pre-emergence (peanuts)	Up to 1.8 L/ha	Queensland dryland	No timing restrictions
		Cape York	No timing restrictions
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	No timing restrictions
		Burdekin	No timing restrictions
		Mackay/Whitsunday	No timing restrictions
		Mary/Burnett	No timing restrictions
		SE Queensland	October and January
Broadcast spray, post-sowing pre-emergence (peanuts)	Up to 3.6 L/ha	Queensland dryland	June to August
		Cape York	October and November
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	October and December
		Burdekin	September to October
		Mackay/Whitsunday	August to December
		Mary/Burnett	April and January
		SE Queensland	Use not supported

Table 2: Application and timing restrictions for application to pastures					
DO NOT apply above maximum rate (L/ha) below OR label rate, whichever is LOWEST					
Pastures (prior to sowing, conservation tillage)	State	Summer	Autumn	Winter	Spring
	Queensland & NT	5.1	5.1	5.1	5.1
	New South Wales & ACT	5.1	5.1	5.1	5.1
	Victoria	0.6	1.7	5.1	1.7
	Tasmania	0.6	1.3	3.6	1.7
	South Australia	1.2	1.7	5.1	3.6
	Western Australia	1.7	3.6	5.1	3.6
Pastures (established)	State	Summer	Autumn	Winter	Spring
	Queensland & NT	7.2	7.2	7.2	7.2
	New South Wales & ACT	7.2	7.2	7.2	7.2
	Victoria	1.0	1.9	7.2	3.6
	Tasmania	0.7	1.7	5.1	3.2
	South Australia	1.4	3.2	7.2	5.1
	Western Australia	3.6	5.1	7.2	5.1

Table 3: Timing restrictions for spraying SUGARCANE		
Rate (L/ha)	Region	Timing Restriction
		DO NOT APPLY DURING THE MONTHS
Up to 1.7 L/ha	Wet Tropics	No timing restrictions
	Burdekin	No timing restrictions
	Mackay/Whitsunday	October to November
	Mary/Burnett	October to November
	Northern NSW	No timing restrictions
Up to 3.6 L/ha	Wet Tropics	October to December
	Burdekin	September to October
	Mackay/Whitsunday	August to December
	Mary/Burnett	April to January
	Northern NSW	October to November

Table 4: Application restrictions for TURF		
DO NOT apply above maximum rate (L/Ha) below OR label rate, whichever is LOWEST		
Turf	State	Rate (L/ha)
	Queensland & NT	3.2
	New South Wales & ACT	3.2
	Victoria	2.5
	Tasmania	2.5
	South Australia	2.5
	Western Australia	4.0

If applying to golf courses in Tasmania, DO NOT apply to fairways adjacent to natural water bodies.

Table 5: Risk mitigation measures for Dryland cropping, pre-emergent uses	
Situation	Risk mitigation measures
Dryland cropping, Preparatory spray	Only apply in no-till farming systems (Tasmania, South Australia)
Winter cereals, preemergence uses	Only apply in no-till farming systems (Tasmania, South Australia, Western Australia)
Summer cereals, preemergent uses	Only apply in no-till farming systems (Tasmania, South Australia)

SPRAY DRIFT RESTRAINTS

DO NOT apply by a vertical sprayer

Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an **unacceptable impact** to **native vegetation, agricultural crops, landscaped gardens** and **aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. The **buffer zones** in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.

DO NOT apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

BOOM SPRAYERS

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a **VERY COARSE spray droplet size category**
- Minimum distances between the **application site** and downwind **sensitive areas**. (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

BUFFER ZONES FOR BOOM SPRAYERS:

Application rate (/ha)	Boom Height above target canopy	Mandatory buffer zones (distance given in metres)				
		Bystander Areas	Natural Aquatic Areas	Pollinator Areas	Vegetation Areas	Livestock Areas
Up to 0.5 L (336 g ae/ha)	0.5m or lower	0	0	0	0	0
	1.0m or lower		30		25	
Up to 1.8 L (1150 g ae/ha)	0.5m or lower		30		25	
	1.0m or lower		60		60	
Up to 3 L (1875 g ae/ha)	0.5m or lower		35		35	
	1.0m or lower		90		90	
Up to 4.4 L (2750 g ae/ha)	0.5m or lower		45		45	
	1.0m or lower		130		130	
Up to 5.3 L (3300 g ae/ha)	0.5m or lower		55		50	
	1.0m or lower		160		160	

AIRCRAFT

DO NOT apply by aircraft unless the following requirements are met:

- Spray droplets not smaller than a VERY COARSE (VC) **spray droplet size** category.
- For maximum release heights above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see the table titled 'Buffer zones for aircraft in the 'Mandatory buffer zones' section below) are observed.

BUFFER ZONES FOR AIRCRAFT:

Application rate (/ha)	Aircraft Type	Mandatory buffer zones (distance given in metres)				
		Bystander Areas	Natural Aquatic Areas	Pollinator Areas	Vegetation Areas	Livestock Areas
Up to 0.9 L (560 g ae/ha)	Fixed Wing	0	120	0	120	0
	Helicopter		85		80	
Up to 1.5 L (950 g ae/ha)	Fixed Wing		170		160	
	Helicopter		120		120	
Up to 1.8 L (1120 g ae/ha)	Fixed Wing		190		180	
	Helicopter		130		130	
Up to 3.6 L (2250 g ae/ha)	Fixed Wing		325		300	
	Helicopter		200		200	

1. FIELD CROPS

CROP	TARGET WEEDS	STATE	RATE L/ha)	CRITICAL COMMENTS	USAGE RESTRICTIONS
Fallow, Stubble Spray Prior to direct drilling or sowing – Winter Cereals, Maize, Sweet Corn, Grain legumes (peanuts Qld, NT only) and Canola.	Refer Weed Table	All States	225 mL- 1.7 L	Observe plant back periods given in the table in this leaflet. Can be mixed with chlorsulfuron or paraquat where grasses are present. Select an appropriate rate from the weed table. For Skeleton Weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.	USAGE RESTRICTIONS APPLY: See Table 1: Timing restrictions for spraying peanuts and Table 5: Risk mitigation measures for Dryland cropping, pre-emergent uses.
Wheat	Refer Weed Table	NSW, SA, ACT only	320 mL - 1.7L	Apply after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller (NSW, SA only)	
		Vic Only	225 mL - 1.4L	Apply from tillering to boot stage (Vic only)	
		Qld Only	560 mL - 1.4L	Apply from mid tillering to before the boot stage (Qld only)	
		Tas Only	800 mL - 1.4L	Apply at 5 leaf to fully tillered (Tas only)	
Barley		NSW, ACT, SA only	320 mL - 1.4L		
		Vic Only	225 mL - 1.4L		
		Qld Only	560 mL - 1.4L		
		Tas Only	800 mL - 1.4L		
Cereal Rye, Triticale		NSW, SA, ACT only	320 mL - 1.4L		
Oats		Vic Only	225 mL - 1.4L		
		NSW, ACT, SA only	320 - 800 mL		
		Vic only	225 - 800 mL		
Cereals: Wheat, Oats, Barley	Cape Tulip Dock, Flatweed, Saffron Thistle Indian Hedge Mustard, London Rocket, Lupin, Matricans, Rapistrum, Wild Radish Wild Turnip	WA only	640 mL - 1.3 L 1.1 L 800 mL 640 mL	Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33) Apply after the 6 leaf stage (Z 16) for Cranbrook, Jacup, Aroona and Spear Wheat and Mortlock Oats to avoid possible damage. DO NOT spray if lucerne is present. WEED STAGE: 10-15 cm. Docks should be sprayed before 5 leaf stage. Cape Tulip – use low rate for cormils only.	

	Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild Turnip		200 mL/ha Plus 500 mL/ha Diuron (500 g/L)	Apply when crop has 4-5 leaves and most weeds have germinated and are in the 2-5 leaf stage. Crop and weeds should be dry at the time of application. Some temporary yellowing of the crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT spray undersown medics. .	
Maize, Sweet Corn	Refer Weed Table	NSW, SA, ACT only	560 mL - 1.1 L/ha	Apply when crop is 10 - 20 cm high and secondary roots are developing for an over-the-top spray. When crop is between 20 cm high and just before tasselling, spray with dropped nozzles to avoid chemical being sprayed into the whort and on upper leaves.	
		Qld only	560 mL - 900 mL/ha	Apply when crop is 10 - 30 cm high and secondary roots have developed. Use drop nozzles and direct the spray when crop exceeds 30 cm in height and before tasselling.	
		Tas only	800 mL – 1.4 L/ha	Apply when crop is 15-30 cm high. DO NOT spray if the crop is showing signs of stress. Some leaf twisting may occur following application – crop recovers quickly.	
Sorghum		NSW, SA, ACT, Vic only	560 mL - 1.1 L/ha	Apply preferably when crop is at 3 – 6 fully expanded leaf stage but can be sprayed from 2 - 8 leaf stage. From 6 leaf stage onwards to within 2 weeks of flowering, crop can be sprayed with dropped nozzles to avoid chemical being sprayed into the whort and on upper leaves.	
		Qld only (except central Qld)	560 mL - 900 mL/ha	Apply when crop has 4 - 8 fully expanded leaves and secondary roots have developed.	
Millet	Refer Weed Table	NSW, SA, ACT, Vic only	560 mL - 1.1 L	Spray when secondary roots have developed, when fully tillered and before heads start to form at the base of the tillers. Do not use on Panorama Millet or Panicum.	
Maize, Sweet Corn, Saccaline, Broom Millet, Millet	Cape Tulip, Dock, Saffron Thistle, Indian Hedge Mustard, London Rocket, Rapistrum, Lupin, Radish, Wild Turnip	WA only	1.1 L	Spray when crop is 10-30 cm high and secondary roots have developed and before tasselling. Apply as direct spray to weeds.	
				Apply when crop is 12 cm high. DO NOT apply between tassel and dough stage. Avoid spraying when in flower.	
Sugar cane (Q80, Q96, and H56 varieties)	Bellvine	Qld, NSW only	280 mL/100 L water	Apply in spring as a direct spray.	USAGE RESTRICTIONS APPLY: SEE TABLE 3: Timing restrictions for spraying SUGARCANE8
	Morning Glory		560 mL – 1.1 L	Apply in summer using a high clearance tractor.	
	Pink Convolvulus, Star of Bethlehem		1.1 L	Apply in autumn by aircraft or misters.	

Sugar Cane	Bindy Eye (Star Burr), Blue Top, Cobblers Pegs, Fleabanes, Jute, Leucas, Needle Burr, Spear Thistle, Water Primrose, Ipomea Vines, Convolvulus Vines	Qld, NSW only	1.8 L - 3.5 L	Add 60 – 120 mL Spraymate Activator to 100 L of spray mixture. Agitate well. DO NOT use on Q63, Q67, Q80 or Q96 varieties.	
	Chinese Mint, Blue Snakeweed		3.5 L		
Peanuts	Broadleaf Weeds, except Noogoora Burr, Grasses, except Mossman Burr	Qld, NT only	1.8 - 3.6 L	LOWER RATE: Apply as BAND SPRAY as soon as possible after planting in a 55cm band. HIGHER RATE: Apply as OVERALL SPRAY after planting and before crop emergence. Some crop damage may occur if heavy rain falls between application and crop emergence.	USAGE RESTRICTIONS APPLY: See Table 1: timing restrictions for spraying PEANUTS
Fallow	Control of Lucerne	All States	2.4 L	Spray in spring when lucerne is actively growing using a minimum spray volume of 50 L/ha. <ul style="list-style-type: none">• Heavily graze lucerne during winter and early spring to reduce crown and root reserves.• Allow lucerne to regrow to 15-30 cm tall before spraying.• Successful lucerne stand reduction is more likely if >70 mm of rain falls in the 6-8 weeks prior to application.• Add either 0.5% Uptake spray oil or a non-ionic surfactant.• Maximum air temperature should not exceed 30°C.	
Harvest Aid or Salvage Spray -Winter Cereals	Desiccate Broadleaf Weeds	All States	1.2 - 1.7 L	Apply after dough stage.	
Bananas	To destroy banana suckers	Qld only	160 mL/10 L water	Inject at the rate of 15 mL per fully grown plant, 10 mL per medium sizes plant and 5 mL for small suckers.	
			320 mL/100 L water	Allow suckers from corms of treated plants to form broad adult leaves, then spray. Isolated spots may require a second spray.	
Cavendish bananas	Banana suckers	All States	160mL/10L water	Apply by stem injection only. INJECT AT THE RATE OF: 15mLper fully grown plant, 10 mL per medium sizes plant and 5 mL for small suckers.	
Common Stylo forage or seed crops Caribbean Stylo forage or seed crops	Refer to Weeds table	Qld Only	800 mL	Apply post-emergence when weeds are 3 weeks old and crop is at least 3 weeks old.	
			800 mL -1.6 L	Apply post-emergence when crop is 3 weeks old.	

2. PASTURES, NON-AGRICULTURAL AREAS, RIGHTS-OF-WAY AND INDUSTRIAL AREAS

CROP	TARGET WEEDS	STATE	RATE (/ha)	CRITICAL COMMENTS	USAGE RESTRICTIONS
Pastures & Non Agricultural Areas	Refer Weed Table	NSW, Qld, SA, ACT, Tas only	560 mL - 1.7 L	Pasture legumes including lucerne, clovers and medics may be damaged unless well protected by grasses. Spot spraying is preferred	USAGE RESTRICTIONS APPLY: See Table 2: Application and timing restrictions for application to pastures
	Galvanised burr	NSW, ACT only	320 mL/100 L water	Apply to young actively growing weeds. Ensure thorough and even coverage of plants. NOTE: Treated plants need to be burnt to destroy seeds.	
	Amsinckia, Annual Thistles, Bindweed, Caltrop, Capeweed, Docks, Flatweed, Heliotrope, Mustard, Paterson's Curse, Ragwort, Saffron Thistle, Spear Thistle, Three cornered Jack (Doublegee, Spiny Emex), Wild Radish, Wild Turnip.	WA only	1.1 L – 2.4 L	For pastures not containing legumes. Only seedling Docks, Spear Thistle and Saffron Thistle will be controlled. SUMMER WEEDS: Use low rate for seedlings, 1.6 - 2.4 L/ha for larger plants. Stock poisoning may occur when grazed after spraying if large amounts present, particularly Heliotrope. WINTER WEEDS: Use low rate for seedlings, 1.6 - 2.4 L/ha for larger plants. If stock are present use spray grazing rates.	
	Afghan Melons		1.6 L/ha plus 1% crop oil	Spray when plants are actively growing preferably before flowering or vining.	
	Paddy melons		800 mL - 1.1 L		
	Prickly Saltwort (Roly Poly)		1.6 L	Spray when plants are small.	
	Stinkwort		1.6 - 3.2 L Plus Surfactant	Best results are obtained when plants are small. Use high rate on larger plants.	
	Dove weed		3.2 L	Spray after good emergence of seedlings.	
Pastures, Rights of way, Industrial Areas.	Boxthorn, Boneseed, Hawthorn	Vic, SA only	Undiluted	Cut stump: Apply or paint as undiluted CropSure Amigo 625 Herbicide to freshly cut stumps.	
	Groundsel	NSW, Qld, SA, ACT only	320 mL/100 L water	HIGH VOLUME: Thoroughly wet plants	
		Tas only	240 mL/15 L water	CUT STUMP: Swab the cut stump immediately. Apply by a pouring can.	
		Qld, NSW, ACT, SA, Tas only	2.9 – 3.6 L	AERIAL APPLICATION: Spray when Groundsel is actively growing.	
	Lantana	NSW, QLD, SA, ACT only	320 mL/100 L water	Use a coarse spray with sufficient pressure to penetrate canopy and wet stems as well as foliage. Spray at the end of a wet summer (March to May). Defoliation should occur but respraying of new growth will be necessary in following Autumn. Broadcast grass seed and keep stock off following Summer to allow the pasture to establish. Damage may result to pasture legumes.	

	Mother of millions	NSW only	400 mL/100 L water	Handgun only: A thorough coverage of leaves and plantlets is necessary. Use 1000g/L non-ionic surfactant at the rate of 1mL of surfactant per 1 L of mixture.	
	Noogoora Burr, Scarlet pimpernel (seedlings only), Weir Vine (<i>Ipomoea</i>), White Eye (Mexican Clover)	Qld only	160 mL/100 L water	In all cases apply to young, actively growing weeds, ensuring through coverage.	
	Annual & Perennial Pigweed, Artichoke Thistle, Bathurst Burr, Billygoat Weed, Blue Snake Weed, Burr Medic, Clockweed*, Fleabanes, Galvanised Burr, Hemlock, Hoary Cress ^o , Kyalinga Weed (Whisker Grass), Knobweed, Milky Cotton Bushes, Parthenium Weed, Paterson's Curse (Salvation Jane), Saffron Thistle, Star Burr, Thornapple, Variegated Thistle*	Qld only	320 mL/100 L water	In all cases apply to young, actively growing weeds, ensuring through coverage. * Spray rosette stage ° Repeat spraying necessary	
	Rubber Vine		160 mL/10 L water	Apply to freshly cut stump.	
	Sesbania pea		560 mL – 900 mL		
	Water hyacinth		3.5 – 5.3 L	Apply in 2,200 to 3,300 L water/ha	
	Wild Tobacco Tree		240 mL/15 L water	Cut Stump Treatment: Swab the cut stump within one hour of cutting. Apply by a pouring can.	
	Agricultural non-crop areas, commercial and industrial areas, pastures and rights of-way	All States	800 mL/ha + wetter	Boom Spray @ 1500 L/ha spray volume To be applied when plant is green. DO NOT apply more than 2 applications per year with a minimum re-treatment interval of 21days between consecutive applications. This product can be used to create and maintain hospital areas for livestock suffering from Pimelea poisoning. Pimelea may become more palatable after herbicide application; stock should be excluded from herbicide-treated areas until sprayed Pimelea plants are leafless, seedless and obviously dead.	
			55 mL/ 100 L water + wetter	Spot Spray. Thoroughly wet all foliage to the point of run-off. (~1500L/ha spray volume)	

Conservation Tillage - Direct Drilling, Surface Sowing or Fallow maintenance	Charlock, Mustards, Shepherd's Purse, Saffron, Slender, Spear and Variegated Thistles*, Turnip Weed, Wild Radish, Wild Turnip.	All States	560 mL – 1.6 L	Apply to actively growing young plants. Before sowing: Observe plant back periods given in the table on this leaflet. * Warning: Treated plants may become toxic to stock.	
	Clover, Sorrel		1.1 L/ha plus 280 - 400 mL/ha of Dicamba (500g/L)	Apply to actively growing plants in autumn. Do not sow pasture seed for at least 30 days after application.	
Lucerne Stand Reduction Fallow Commencement	Lucerne	All States	2.4 L/Ha	Spray in spring when lucerne is actively growing using a minimum spray volume of 50 L/ha. Heavily graze lucerne in winter and early spring to reduce crown and root reserves. Allow lucerne to regrow to 15 - 30 cm before spraying. Successful lucerne stand reduction is more likely if >70 mm of rain falls in the 6 to 8 weeks prior to application. Add either 0.5% CropSure Takedown Spray Oil or a non-ionic surfactant. Maximum air temperature should not exceed 30°C.	USAGE RESTRICTIONS APPLY: See Table 2: Application and timing restrictions for application to pastures
Pastures - Spray/Graze Techniques	Precaution: An increased quantity of poisonous plants may be eaten by stock using spray-graze e.g., Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended period of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse				
	Amsinckia, Thistles, Capeweed, Doublegee, Mustards, Paterson's Curse, Wild Turnip, Wild Radish, Docks, Geranium, Erodium	SA only	560 mL	Apply from 6 weeks after opening rains in autumn until the end of August. Seven days after spraying stock paddock at 4-5 times normal rate, preferably with sheep. Maintain this level of grazing for 6 weeks or until pasture shows signs of over grazing, then return to normal stocking levels. Use high stocking rates in following spring to prevent weeds from flowering. Repeat treatments may be required for 2-3 years for complete control. Refer to " Precaution " above	USAGE RESTRICTIONS APPLY: See Table 2: Application and timing restrictions for application to pastures
	Annual Thistles, Capeweed, Doublegee (Three Cornered Jack, Spiny Emex), Geranium, Mustards, Paterson's Curse (Salvation Jane), Saffron Thistle, Slender Thistle, Spear Thistle, Wild Turnip,	Tas & Vic only	600 - 800 mL		
	Amsinckia, Annual Thistles, Capeweed, Docks (seedlings only), Doublegee (Three Cornered Jack, Spiny Emex), Mustard, Paterson's Curse (Salvation Jane), Wild Radish, Wild Turnip.	WA only			

	Saffron Thistle, Spear Thistle		1.2 L	Apply to Saffron Thistle at the end of September when plants are running up to flower. Sub-clovers may be damaged at this rate and use is not recommended for all medic pastures.	
	Melons	WA only	1.6 L/ha Plus 1% oil	Heavy stocking on young plants sprayed with 800 mL/ha provides effective control.	
	Docks	Vic only	1.1 L	Apply in September and follow other recommendations above.	
	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse (Salvation Jane), Shepherd's Purse, Saffron, Slender, Spear or Variegated Thistles*, Turnip Weed, Wild Radish, Wild Turnip	NSW & ACT only	280 mL - 1.1 L	Spray actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep (cattle are less effective). Stocking rate should be at least 5 times heavier than normal until weeds have been reduced, but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed shortly before spraying. Other legumes may be affected. * Warning: treated plants may become toxic to stock	

3. LAWNS AND PLANTATIONS

CROP	TARGET WEEDS	STATE	RATE (/ha)	CRITICAL COMMENTS	USAGE RESTRICTIONS
Lawns, Playing fields	Refer to weed table	Qld, NSW, ACT only	1.5mL-3mL/1L of water or 1.6 - 3.2L/ha	Wet foliage thoroughly. DO NOT mow lawn for 1 week before and at least 1 week after application. DO NOT use on Buffalo Grass (WA only).	USAGE RESTRICTIONS APPLY. SEE TABLE 4: Application restrictions for TURF.
		WA only	40mL per 10-15L water/100m ²		
Hardwood and Softwood Plantations	Broadleaf and woody weeds as per product labels including groundsel and Pinus spp. Wildings	All States	Maximum rate 1.6 L/h	Apply a single preplant application and/or a maximum of 2 post plant applications using shielded sprayers within the first 2 years following planting. Apply using aircraft (rotary wing only) or ground based equipment. DO NOT spray over or into watercourses. Products may be mixed with glyphosate for pre-plant spray operations	
Oil Tea Tree	Weeds on label	All States	Apply at a maximum of 960 mL/ha as per label directions	Apply as a shielded spray. Avoid contact with foliage, green stems, exposed non- woody roots, desirable plants and trees as severe injury or destruction may result. Apply following harvest as a blanket spray only after: • All residual tea tree foliage has been removed by mechanical shaving, or by using a burner, • No swollen buds are present on stumps. NOTE that buds can burst 8 days after harvest in summer and • Surface of cut stumps are dry before spraying commences.	
	Purple Top (Verbena bonariensis)		Apply at 960mL/ha plus 720 g ai glyphosate/ha in tank mix.	Apply as a shielded spray. Avoid contact with foliage, green stems, exposed non-woody roots, desirable plants and trees as severe injury or destruction may result.	

4. SPOT SPRAYING

SITUATION/ CROP	TARGET WEEDS	STATE	RATE (/ha)	CRITICAL COMMENTS	USAGE RESTRICTIONS
Spot spraying High Volume Spraying	Refer Weed Table	All States		Add 1/10 th of the rate on the weeds table to 150 litres of water. Each 150 litres of mix will cover 1000 m ² (1/10 th ha) e.g. If rate in weed table is 1.4 L use 140 mL/150 L water.	

MAXIMUM TOLERANCE FOR USE IN WHEAT, BARLEY, OATS AND TRITICALE INCLUDING TOLERANCE FOR UNDERSOWN LEGUMES							
CROP	QLD	NSW/ACT	VICTORIA		SA	WA	TAS
			Early tillering	Tillered boot stage		Tillered to Boot (Z15-35)	Tillered to Boot (Z15-35)
Cereal Rye	-	-	-	-	1.4 L	-	-
Wheat	1.8 L/ha	1.7 L/ha	225 mL/ha	1.4 L/ha	1.7 L/ha	1.7 L/ha	1.7 L/ha
Barley	1.4 L/ha	1.4 L/ha	225 mL/ha	1.4 L/ha	1.4 L/ha	1.4 L/ha	1.4 L/ha
Oats	-	800 mL/ha	225 mL/ha	800 mL/ha	900 mL/ha	1.3 L/ha	-
Triticale	-	1.4 L/ha	225 mL/ha	-	1.4 L/ha	1.7 L/ha	800 mL/ha
Undersown clovers	-	-	225 mL/ha	-	-	680 mL/ha	560 mL/ha
Undersown medics	-	-	225 mL/ha	-	-	Nil	560 mL/ha
Undersown lucerne	-	-	225 mL/ha	-	-	Nil	-

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

Pasture, Cereal Crops - **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

Harvest- **NOT REQUIRED WHEN USED AS DIRECTED**

Fallow, Lucerne - **DO NOT GRAZE, CUT OR CULTIVATE FOR AT LEAST 21 DAYS AFTER SPRAYING.**

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the particular crop or situation indicated under the Directions for Use Table.

In most cases, this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

WEEDS	APPLICATION RATE PER HECTARE						CRITICAL COMMENTS	
	CROP							
	VIC	NSW, ACT	SA	QLD	TAS	WA		
Amaranthus spp.	-	560mL- 1.1L	-	900mL	-	-	-	
Amsinckia	-	-	-	-	-	1.1L	-	
Apple of Peru	-	560mL - 1.1L	-	900mL	-	1.1 L	-	
Bathurst Burr	-	800mL- 1.1L	1.7 - 2.2L	900mL	-	-	800mL - 1.1L Not SA	
Bellvine	-	1.8L	-	1.8L	-	-	-	
Billygoat Weed	-	3.4L	-	3.4L	-	-	-	
Bindweed	-	-	-	-	-	1.1L	-	
Blackberry Nightshade	-	560mL - 1.1L	-	900mL	-	-	-	
Blackeyed Susan	-	1.8L	-	1.8L	-	-	-	
Blue Snakeweed	-	1.8L	-	1.8L	-	-	-	
California Burr	-	800mL- 1.1L	-	900mL	-	-	800mL-1.1L not SA	
Cape Tulip	-	-	-	-	-	640mL -1.3L	-	
Capeweed	1.4L	-	1.7L	-	1.4L	1.1L	1.7-2.8L	
Caltrop	-	800mL- 1.7L	-	900mL	-	1.1L	-	
Castor Oil Plant	-	3.4L	-	3.4L	-	-	-	
Charlock	560mL- 800mL	560mL- 1.1L	560mL	-	1.4L	-	800mL-1.1L	
Clover	-	1.2L	-	-	-	-	-	
Cobbler's Pegs	-	1.8L	-	1.8L	-	-	-	
Common Ice Plant	-	-	1.1L	-	-	-	-	
Common Sida	-	1.8L	-	1.8L	-	-	-	
Common Sowthistle	-	1.8L	-	1.8L	1.4L	-	-	
Docks	1.4	-	1.4L	1.4L	1.4L	1.1L	3.2L SA only	
Doveweed	-	-	-	-	-	1.1L	-	
Fat Hen	-	560mL- 1.7L	-	900mL	1.4L	-	-	
Flannel Weed	-	1.8L	-	1.8L	-	-	-	
Flat Weed	-	-	-	-	-	1.1L	-	
Fumitory – red	-	-	1.7L	-	-	-	-	
Fumitory – white	800mL	-	560mL	-	-	-	-	
Heliotrope	-	-	-	-	-	1.1L	-	
Hexham Scent or Meliolotus	1.4L	-	1.1L	1.4L	-	-	1.1-1.7L	
Hoary Cress	900mL- 1.4L	1.1-1.7L	1.4L	1.4L	-	-	1.4-1.7L	
Hogweed/ Wireweed	1.4L	-	-	1.4L	-	-	-	
							Spray at multiple leaf stage (Vic) Spray at seedling and young plant stage (Qld).	

WEEDS	APPLICATION RATE PER HECTARE						CRITICAL COMMENTS	
	CROP							
	VIC	NSW, ACT	SA	QLD	TAS	WA		
Horehound	-	-	1.4L	-	-	-	2.2-3.2L SA only	Spray seedlings.
Indian Hedge Mustard	-	-	-	-	1.4L	1.1L	-	
Khaki Weed	-	-	-	-	-	-	1.1-2.2L not SA	Spray seedlings only
Knobweed	-	-	-	1.8-3.4L	-	-	-	Lower rate for seedlings; higher rate for later stages.
Lincoln Weed	-	-	1.7L	-	-	-	-	Spray early rosettes
London Rocket	-	-	-	-	-	1.1L	-	
Lupins	-	800mL- 1.7L	-	-	-	1.1L	-	
Melons – Camel, Paddy	-	560mL- 1.1L	-	-	-	-	-	
Mexican Poppy	-	-	-	1.4L	-	-	-	Spray seedlings - plants become more resistant with age.
Mintweed	-	1.1L	-	900mL	-	-	-	Spray seedlings - resistant in later stages.
Morning Glory	-	1.8L	-	1.8L	-	-	-	Spray at seedling to flowering stage.
Mustards	225- 560mL	560ml- 1.1L	560ml- 1.4L	900mL	-	800mL	560ml-1.1L	Spray at 2-4 leaf up to rosette stage.
Needle Burr	-	1.8L	-	1.8L	-	-	-	Apply at pre-flowering preferably young stages.
New Zealand Spinach	-	1.1-1.7L	-	-	-	-	-	
Noogoora Burr	-	800mL- 1.1L	-	900mL	-	-	800mL-1.1L not SA	Spray seedlings only
Paterson's Curse	-	1.1-1.7L	-	1.4L	-	1.3L	1.7-2.2L	Spray rosettes at or before plants have 10 leaves. Later stages harder to kill.
Pinkburr (Pink Flowered Burr)	-	1.8L	-	1.8L	-	-	-	Spray seedlings or young stages only.
Potato Weed	-	560mL- 1.1L	-	900mL	-	-	-	
Purpletop	-	3.4L	-	3.4L	-	-	-	Spray at young stage.
Radish	-	-	-	-	-	1.1L	-	
Ragwort	-	-	-	-	3.8L	1.1L	-	Spray up to early rosette stage (Tas only).
Rapeseed	-	800mL- 1.7L	-	-	-	-	-	
Rapistrum	-	-	-	-	-	1.1L	-	
Rough Poppy	-	1.1L	-	-	-	-	-	
Safflower	-	560mL- 1.1L	-	-	-	-	-	
Shepherds Purse	-	1.1 – 1.7L	-	-	1.4L	-	800mL-1.1L	Spray young rosettes
Siratro (Purple Bean)	-	1.8L	-	1.8L	-	-	-	Spray seedlings at young stages only.
Skeleton Weed	1.4L	1.1-1.7L	1.4L	-	-	-	-	Spray rosettes before aerial growth commences.
Sorrel	1.4L	1.7L	1.4L	-	-	-	-	Only moderately susceptible
Speedwell - Ivy leaf	-	-	1.1L	-	-	-	-	
Spinyhead Sida	-	1.8L	-	1.8L	-	-	-	Spray seedling or young stages only.
Starburr	-	1.8L	-	1.8L	-	-	-	Spray before seeding, advanced stages susceptible.
Spiny Emex	-	-	-	1.4L	-	-	-	Only young plants are susceptible.
Star of Bethlehem (Cupid's Flower)	-	-	-	1.8L	-	-	-	Spray before seeding, advanced stages susceptible.
Stinkwort	-	800mL- 1.4L	-	-	-	-	-	
Storksbill/Erodium	-	-	-	-	1.4L	-	1.6-3.2L	Spray seedlings to young rosettes.
Sunflower (seedlings)	1.4L	560mL- 1.4L	-	900mL	-	-	-	

WEEDS	APPLICATION RATE PER HECTARE						CRITICAL COMMENTS	
	CROP							
	VIC	NSW, ACT	SA	QLD	TAS	WA		
Thistles								
- Annual	-	-	-	-	-	1.1L	-	
- Californian	-	-	-	-	2.7L	-	3.2-3.8L	
- Saffron	1.1L	560mL- 1.7L	1.4L	1.4L	950mL	1.4L	1.1-1.7L	
- Slender/ Shore	-	800mL- 1.7L	-	-	1.4L	1.4L	1.1L	
- Soldier	1.4L	-	-	-	-	-	1.1-1.6L not NSW, Tas	
- Spear	560mL	-	-	-	1.4L	-	1.1-1.6L	
- Star	-	-	-	-	-	-	1.6-3.2L SA only	
- Variegated	-	560mL- 1.7L	-	900mL	1.4L	-	1.1-1.7L	
Thornapple	-	800mL- 1.1L	-	-	-	-	1.6-2.4 not SA	
Tridax (Tridax Daisy)	-	1.8L	-	1.8L	-	-	Spray seedling or young stages only.	
Turnip Weed/Rapistrum	-	560mL- 1.1L	-	560mL	-	800mL	560ml-1.1L	
Vetches/Tares	1.4L	-	1.1L	-	-	-	Spray at multiple leaf stage.	
Wards Weed	-	-	1.1L	-	-	-	-	
Wild Cabbage	1.4L	-	-	-	-	-	Spray multiple leaves.	
Wild Poppy	560mL	-	-	-	-	-	Spray rosettes.	
Wild Radish	1.4L	1.4-1.7L	1.4L	900mL	1.4L	800mL	800mL-1.1L	
Wild Turnip	225- 560mL	560mL- 1.1L	320mL	-	1.4L	640ml	560mL-1.1L	
							Spray 2-4 leaf up to rosette stage.	

PLANT BACK PERIODS (DAYS) FOR CropSure Amigo 625 Herbicide

CROP/RATES	UP TO 560 mL/ha	560mL – 1.1 L/ha	1.1 – 1.7 L/ha
Balansa Clover	7	7	10
Barley %	1	1	3
Chickpea #	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medic	7	7	10
Narbon Beans	7	7	10
Navybeans	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White Clover	7	7	10

IMPORTANT:

WHEN APPLIED TO DRY SOILS AT LEAST 15 mm (1/2 inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

- % In Queensland, no rainfall is required to fall prior to commencement of the Plant Back Period for wheat, barley and triticale.
- # In Queensland, planting of canola/rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15 mm.
- @ In Central Queensland, when using 800mL/ha or less of CropSure Amigo 625 Herbicide, the Plant back Period for sorghum and sunflower is 1 day irrespective of rainfall.
- + In WA the Plant back Period for Lupins at all rates is 28 days.

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

Mode of action

This product is readily absorbed through plant roots. The chemical mimics the action of auxins causing over stimulation of growth and rupture of cells and eventual death of susceptible weeds.

Mixing

CropSure Amigo 625 Herbicide mixes readily in water. Part fill spray tank with water. Add the required amount of CropSure Amigo 625 Herbicide and fill the rest of the tank. Maintain agitation.

APPLICATION INFORMATION:

- BOOM SPRAYING - Use 30 – 120 L/ha of water.
- AERIAL SPRAYING - Use 10 – 90 L/ha of water

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy water or 1% solution of Ammonia followed by several clean water rinses

COMPATIBILITY

CropSure Amigo 625 Herbicide is compatible in tank mixes with 500 g/L Dicamba, 500g/L Diuron Flowable, 900 g/kg Diuron, Paraquat, 900 g/kg Atrazine, 500 g/L Atrazine Flowable, Paraquat & Diquat mix, Methabenzthiazuron, 2,2-DPA, Picloram and Chlorsulfuron

RESISTANT WEEDS WARNING:**GROUP 4 HERBICIDE**

CropSure Amigo 625 Herbicide is a member of the Phenoxy group of herbicides. CropSure Amigo 625 Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management, CropSure Amigo 625 Herbicide is a GROUP 4 HERBICIDE. Some naturally occurring weed biotypes resistant to CropSure Amigo 625 Herbicide and other GROUP 4 HERBICIDE may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CropSure Amigo 625 Herbicide or other GROUP 4 HERBICIDES.

Since the occurrence of resistant weeds is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Amigo 625 Herbicide to control resistant weeds.

RE-ENTRY PERIOD

If re-entering treated areas before the spray has dried, workers should wear overalls, elbow-length gloves and water-resistant footwear. Clothing must be laundered after each day's use.

DO NOT hand harvest sugar cane for at least 1 day after application.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray cereals if lucerne is present.

DO NOT spray crops or weed outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants, crops, cropped lands or pastures.

Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF HONEYBEES AND OTHER INSECT POLLINATORS

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. **DO NOT** contaminate streams, rivers or watercourses with the chemical or used container.

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

For Mini Bulk Returnable Containers

Store the original sealed container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the MicroMatic valve or the security seal. DO NOT contaminate the container with water or any other foreign matter. After each use of the product ensure the MicroMatic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the container have been used, please return the empty container to the point of purchase. The container remains the property of CropSure Pty Ltd.

SAFETY DIRECTIONS:

Poisonous if absorbed by skin contact, inhaled or swallowed. Corrosive to the eyes and skin. Will irritate the nose and throat. Avoid contact with the eyes and skin. Do not inhale vapour or spray mist.

When opening the container and preparing spray or using undiluted concentrate, wear chemical resistant waterproof clothing over a layer of normal clothing and a washable hat, elbow-length chemical resistant gloves, impervious footwear and full face piece respirator with organic vapour/gas cartridge or canister.

If applying by boom spray equipment with enclosed operator's cab and air filtration or aerial spraying equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.

If applying by boom spray equipment with open operator's cab or hand-held spray equipment wear chemical resistant waterproof clothing over a layer of normal clothing and a washable hat, elbow-length chemical resistant gloves and full face piece respirator with organic vapour/gas cartridge or canister.

If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves (and) respirator and if rubber wash with detergent and warm water contaminated clothing.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**