

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure **Metribuzz 750WG** **Herbicide**

ACTIVE CONSTITUENT: 750g/kg METRIBUZIN

GROUP 5 HERBICIDE

**For Selective Weed Control in Cereals, Pastures and other Crops,
including Vegetables, as Indicated in the Directions for Use**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING
OR USING THIS PRODUCT**

Contents: 10 kg

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury containers in a local landfill. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) that can be obtained from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

24HR Emergency Response:
AU +61 1800 951 288
NZ +64 800 700 112

APVMA Approval No.: 90405/128772

Date of Manufacture:

Harmful if swallowed. Very toxic to aquatic life with long lasting effects. Avoid breathing dusts. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Brush off loose particles from skin. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. In case of fire: Use carbon dioxide, dry chemical, foam, water fog, to extinguish. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure **Metribuzz 750WG** **Herbicide**

ACTIVE CONSTITUENT: 750g/kg METRIBUZIN

GROUP

5

HERBICIDE

**For Selective Weed Control in Cereals, Pastures and other Crops,
including Vegetables, as Indicated in the Directions for Use**

**IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING
THIS PRODUCT**

APVMA Approval No.: 90405/128772

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply to crops that are stressed by any cause (such as severe weather conditions, waterlogging, drought, excessive soil acidity or alkalinity, poor nutrient status, disease, nematode or insect damage) as crop injury may result.

Crop / Situation	Weeds Controlled	State	Rate / ha	Critical Comments
Barley	Capeweed, Cat's Ear, Charlock, Chickweed, Common Cotula, Corn Gromwell (Sheepweed, White Ironweed), Corn Spurry, Creeping Speedwell, Crown Beard (Dogweed), Deadnettle (Henbit Lamium spp.) Dock seedlings, Fat Hen, Field Madder, Fumitory, Hare's Ear (Treacle Mustard), Heliotrope, Hogweed (Wireweed), Horehound, Indian Hedge Mustard, Lesser Swinecress, Mouse-ear Chickweed, Mountain Sorrel, (Purple Calandrinia), Rough Poppy, Scarlet Pimpernel, Scrub Nettle (Stinging Nettle), Shepherds Purse, Sorrel seedlings, Sowthistle, Spiny Emex (Doublegee, Three cornered Jack(s)), Storksbill, Toad Rush, Wild Radish, Wild Turnip, Winter Grass, Yellow Burrweed	Vic & Tas only	180g (Light sandy soils, i.e., sandy loams to loamy sands) 280g (Medium soils, loams, silt plus clay 40-60%) 380g (Heavy soils, clay loams, silt plus clay over 60%)	Apply to moist soils while weeds are young and actively growing and barley plants are at the 5-leaf stage. DO NOT use on Barley grown on alkaline light sandy soils with pH over 7. DO NOT use on Barley grown on light sandy soils (Tas).
Barley	Buxbaum's Speedwell, Powell's Amaranth, Stagger weed, Sun Spurge, Great Brome	Tas only		
Barley All varieties excluding Morrell	Capeweed, Common Cotula, Doublegee, Fumitory, Indian Hedge Mustard, Toadrush, Wild Radish, Wild Turnip, Wireweed. Brome grass (Bromus diandrus & B. rigidus) Annual Ryegrass, Barley grass	WA only Sandy - sandy loam soils	135-150g plus trifluralin 480 at 1.0L/ha and incorporated by sowing (IBS)	DO NOT apply to Morell barley. Use the lower rate in lighter soils, and the higher rate in heavier soils. Apply to moist soil for best efficacy Seed below herbicide band at 5cm. Reduced control may occur under high weed burdens (>100/m ²). Ensure CropSure Metribuzz 750WG Herbicide is fully dispersed in the boom spray tank prior to the addition of Trifluralin 480. Refer to the application section of this and the trifluralin label for complete instructions on product use.
Chick peas, Faba Beans, Lentils, Vetch	Refer to the barley section	SA, VIC & WA only	180 g (Light sandy soils, i.e., sandy loams to loamy sands)	Crop: ONLY apply post-sowing, pre-emergence at least 5 cm deep. DO NOT apply post-emergence, as crop injury will occur. Weeds: Weeds should be from pre-emergence to 3-leaf stage except Wireweed (Hogweed) that should not be beyond the cotyledon stage. Best results for grass

Crop / Situation	Weeds Controlled	State	Rate / ha	Critical Comments
			280g (Medium soils, loams, silt plus clay 40-60%) 380g (Heavy soils, clay loams, silt plus clay over 60%)	and Wireweed control are obtained when rain or irrigation follows within a few days of spraying.
Lupins	Capeweed, Doublegee, Wild Radish	WA only	100-150g plus 100mL diflufenican	For use on varieties Gunurru, Meritt, Myallie and Kayla only. DO NOT use on Danja, Kiev mutant or Albus lupins. Suppression of larger Doublegee only. Wild Radish weed size up to 250mm only. Apply to the actively growing lupins from the 3-4 leaf stage onwards. Use higher rates for heavy weed burden and larger weed size. Apply before weeds are covered by crop canopy. DO NOT apply wetting agents or spray oils with this mixture. Some transient leaf burn may occur after application, but this will not affect yield. Best results are obtained if good soil moisture exists at, and is received soon after application. Results may be less effective south of the Great Eastern Highway. Refer to critical comments section on the diflufenican label.
Peas (Field and Green) (Not Snow Peas)	Refer to Barley section plus: Buchan Weed, Ball Mustard, Fog Grass (SA only), Dwarf Nettle (Tas only) Suppression only- Sub clover and Docks (Broadleaved Dock – Tas only), Wild Oats	NSW, Vic, SA & Tas only	180g (Light sandy soils, i.e., sandy loams to loamy sands) 280g (Medium soils, loams, silt plus clay 40-60%) 380g (Heavy soils, clay loams, silt plus clay over 60%) 470g (Kooweeerup swamp and volcanic soils)	Crop: Sow crop 3-4cm deep. Apply after rain when crop is pre-emergence to 3-node stage (Refer to diagram below for timing of application to crop). For post-sowing pre-emergence application allow soil to settle for 2-3 days after sowing before spraying. For post-emergence apply to moist soil within 6 weeks of sowing while weeds are young and actively growing. DO NOT spray crops affected with collar or stem rot. (Diagram 1-3 nodes). Weeds: Weeds should be from pre-emergence to 3-leaf stage except Wireweed (Hogweed) that should not be beyond the cotyledon stage. Best results for grass and Wireweed control are obtained when rain or irrigation follows within a few days of spraying.

Crop / Situation	Weeds Controlled	State	Rate / ha	Critical Comments
Peas (Field and Green) (Not Snow Peas) continued	Refer to Barley section plus: Buchan Weed, Ball Mustard, Fog Grass (SA only), Dwarf Nettle (Tas only).	QLD only	470g 700g	Apply to medium soils Apply to soils high in clay or organic matter content when grass weeds are a problem.
Green Peas (Post emergence) (NOT Snow Peas)	Suppression only- Sub clover and Docks (Broadleaved Dock – Tas only), Wild Oats	WA only	370-470g	Refer to Critical Comments for Peas (Field and Green)
Potatoes		ACT, NSW, VIC, SA, Tas & WA only	470g 700g	Apply after rain or irrigation when crop has no more than 6 leaves or is no more than 10 cm high (many plants may be smaller). CROPS HILLED OR MOLDED UP – Hill or mold up as the crop emerges. Apply 1-4 weeks later as soon as weed seedlings have emerged. If surface crusting is not a problem no further cultivation should be necessary. CROPS NOT HILLED AT OR SOON AFTER PLANTING - Spray when weed seedlings have emerged. No cultivation should be necessary until crop is ready for final hilling. Additional cultivation may be necessary when surface crusting is a problem.
		Vic, Bellarine Peninsula only	280-370g	Where summer grasses are a problem.
		Qld only (Winter crop)	470g	Apply to winter crops on light soils only.
		South Qld only (Summer crop)	470g - 950 g	Apply at first emergence of potatoes to control broadleaf weeds.
		Atherton Tablelands (QLD only)	950g	Where summer grasses such as <i>Echinochloa crus-gali</i> , <i>Eleusine</i> , <i>Digitaria</i> and <i>Panicum spp.</i> are a problem.
				Apply to summer crops at first emergence of potatoes to control broadleaved weeds and more troublesome grasses such as <i>Eleusine</i> , <i>Digitaria</i> and <i>Panicum spp.</i>
Irrigated Soybeans	Apple of Peru, Cobbler's Pegs, Pigweed, Noogoora Burr, Common Thornapple, Summer Grass, Caltrop	ACT, NSW, NT only	500g	Apply pre-plant incorporated. Where grass weeds other than summer grass are a problem apply in combination with trifluralin 480. (Refer to label for directions for use and incorporation techniques). Use only on soils classified as silty clay-heavy clay.
Irrigated Soybeans (Variety Davis only)	Refer to weeds listed for soybeans plus <i>Anoda</i> weed, Awnless Barnyard grass. Suppression only of Bellvine	QLD, NT only	750g	Apply pre-plant incorporated. Where grass weeds other than summer grass are a problem apply in combination with trifluralin 480. (Refer to label for directions for use and incorporation techniques). Use only on soils classified as silty clay-heavy clay.
Tomatoes Inter-row spaces of furrow	Refer to Peas section plus: Barnyard grass, Summer grass, Crowsfoot grass, Blackberry Nightshade, Pigweed, Black Pigweed, Bull's Head, Green Summer Grass		950g or 9.5g/100 m of row	Apply in a 1 metre band centred on the interrow either during or shortly after the formation of the final furrow. Apply to soil moist at the surface. DO NOT exceed spraying pressures of 200 kPa, as excessive drift will occur.

Crop / Situation	Weeds Controlled	State	Rate / ha	Critical Comments
irrigated tomatoes				
Wheat (Blade only)	Capeweed, Common Cotula, Doublegee, Fumitory, Indian Hedge Mustard, Toad Rush, Wild Turnip, Wild Radish, Wireweed. Brome Grass (Bromus diandrus and B. rigidus) Annual Ryegrass, Barley Grass (refer to critical comments)	WA- Great northern region only Sandy- sandy loam soils	200g and incorporated by sowing (IBS) 200g/ha plus a trifluralin 480 at 1L/ha and incorporated by sowing (IBS)	CropSure Metribuzz 750WG Herbicide has increased crop safety and gives more reliable weed control when incorporated by sowing and trailing harrows. Reduced grass control may occur under high weed burdens (>100/m ²). To achieve good control of Annual Ryegrass and Barley grass add trifluralin 480 at 1L/ha. Ensure CropSure Metribuzz 750WG Herbicide is fully dispersed in the boom spray tank prior to the addition of trifluralin 480. Refer to the application section of this and the trifluralin label for complete instructions on product use.
Wheat and Barley	Toad Rush	Vic, ACT, NSW, SA only	100g	Toad rush should be sprayed at the 2-4 leaf stage for optimum control.
Oats		Vic only		Spray when soil moisture is plentiful and soil is moist to the surface. Take advantage of dew on the soil surface.
White Lupins	Buchan Weed, Capeweed, Charlock, Chickweed, Deadnettle (Lamium spp.), Fat Hen, Fumitory (Common and Wall Fumitory- Tas), Wireweed (Hogweed), Lesser Swinecress, Purple Calandrinia, Scarlet Pimpernel, Spiny Emex, Corn Spurry (Spurry-Tas), Stinging Nettle (Dwarf Nettle-Tas), Toad Rush, Winter grass, Heliotrope, Wild Oats, Wild Turnip, Wild Radish. Suppression only: Sub clover and docks (Broadleaved dock-Tas).	Tas only	400g	Spring application – Apply after sowing but before crop emerges.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD**TOMATOES: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION****BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION****OTHER CROPS: NOT REQUIRED IF USED AS DIRECTED****GENERAL INSTRUCTIONS**

CropSure Metribuzz 750WG Herbicide is non-corrosive to equipment, non-flammable and non-volatile. The degree of weed control and duration of effect will vary with the amount of herbicide applied, soil texture, rainfall and other conditions.

RESISTANT WEEDS WARNING

GROUP	5	HERBICIDE
-------	---	-----------

CropSure Metribuzz 750WG Herbicide is a member of the Triazines group of herbicides. CropSure Metribuzz 750WG Herbicide has the inhibitors of photosynthesis at Photosystem II mode of action. For weed resistance management CropSure Metribuzz 750WG Herbicide is a Group 5 Herbicide.

Some naturally- occurring weed biotypes resistant to CropSure Metribuzz 750WG Herbicide and other Group 5 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CropSure Metribuzz 750WG Herbicide or other Group 5 herbicides.

Annual Ryegrass biotypes resistant to diclofop-methyl and other ryegrass specific herbicides may be resistant to CropSure Metribuzz 750WG Herbicide. CropSure Pty Ltd recommends resistance testing of the population to ensure susceptibility prior to use.

Since the occurrence of resistant weeds is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Metribuzz 750WG Herbicide to control resistant weeds. Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the survivors.

DO NOT make more than one application of a Group 5 Herbicide to a crop in the same season, either pre-sowing incorporated by sowing or post crop and weed emergence. If the user suspects that Group 5 Herbicide resistant weeds are present, CropSure Metribuzz 750WG Herbicide or other Group 5 Herbicides should not be used.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

CROP SAFETY

DO NOT apply to crops that are stressed by any cause (such as severe weather conditions, waterlogging, drought, excessive soil acidity or alkalinity, poor nutrient status, disease, nematode or insect damage) as crop injury may result.

Barley:	DO NOT apply to crops grown on sandy soils or soils low in organic matter.
Peas:	DO NOT apply to Laura, Pennant or White Brunswick varieties as phytotoxicity may occur.
(field & green)	DO NOT apply any later than the one node stage to Alma and Dinkum peas.
Potatoes	DO NOT spray pea crops in soil s or areas where collar or stem rot is likely to occur.
	DO NOT apply to crops grown on sandy soils as injury to the crops may result, appearing as chlorosis or minor necrosis.
	DO NOT apply to Kurell potatoes as phytotoxicity may occur.
	DO NOT apply to Alantic and Pontiac varieties – WA only.
	DO NOT use on dry-grown, non-irrigated crops.
Soybeans:	DO NOT use on varieties Cannapolis, Hill and Semstar as crop injury may occur.
	DO NOT use on soils with a calcareous surface layer as crop injury may occur.
	DO NOT apply over atrazine residues as crop injury may result.
	DO NOT apply in combination with soil applied organophosphate insecticides as crop injury may result.
	DO NOT apply to soils with less than 0.5% organic matter as crop injury may result.
	DO NOT apply to dryland soybeans.
	DO NOT apply to soils of pH greater than 8 that are prone to waterlogging as crop injury may occur.
Lupins	DO NOT apply to varieties Danja, Kiev Mutant or Albus Lupins.
	DO NOT apply to Lupin crops that have received 2.5 litres or more of Simazine pre-seeding, or lupin crops that are showing symptoms of Simazine or Atrazine damage.

MIXING AND APPLICATION COMPATIBILITY

CropSure Metribuzz 750WG Herbicide is compatible with the herbicides: chlorsulfuron, metsulfuron, glyphosate, pendimethalin, propargite, diflufenican and trifluralin 480.

MIXING**General**

CropSure Metribuzz 750WG Herbicide is to be mixed with water and applied as a spray. Before spraying calibrate equipment to determine the quantity of water necessary to uniformly cover the measured area to be treated. Partially fill the tank with water. Measure the amount of herbicide required for the area to be sprayed, adding this amount of CropSure Metribuzz 750WG Herbicide to the spray tank with the agitation system engaged. Top up to the correct volume with water.

THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.

In tank mixes CropSure Metribuzz 750WG Herbicide must be in suspension before adding the companion herbicide or liquid fertiliser.

Pre-planting Incorporation Treatment:

CropSure Metribuzz 750WG Herbicide plus trifluralin 480:

Add CropSure Metribuzz 750WG Herbicide to water in spray tank with agitator running as above. Maintain agitation then add trifluralin 480 at the recommended rates (Refer to trifluralin 480 label).

APPLICATION**General**

Best results are obtained if treatment is made to moist soil and moisture is supplied by rainfall or sprinkler irrigation (6-12mm) within 2 weeks after application. If heavy rains occur soon after application, injury to the crop may result.

Aerial Application - (Qld only)

Peas-apply in a minimum of 20 litres water per hectare.

Potatoes - apply in a minimum of 50 litres water per hectare. Avoid overlapping of spray swath. Avoid spraying in still conditions and in wind likely to cause drift.

Pre-emergence and Post-emergence (to crop)

CropSure Metribuzz 750WG Herbicide should not be applied when weeds are in the young seedling stage (no more than 3 true leaves). Under dry conditions there is no root uptake of CropSure Metribuzz 750WG Herbicide and results against grass weeds and Wireweed (Hogweed) will be poor. Except for small areas, use fixed boom power sprayers set at 200-300kPa (for tomatoes do not exceed 200kPa) and fitted with fan nozzles. Apply a minimum of 80L-100L prepared spray/ha. Openings in screens should be equal to or larger than 50 mesh. Agitate by mechanical or hydraulic means in spray tank. When spraying avoid overlapping. Shut off spray booms whilst starting turning, slowing or stopping otherwise crop injury may result.

Pre-plant incorporated - Soybeans only

CropSure Metribuzz 750WG Herbicide alone or in combination with Trifluralin 480

Spray anytime from 4 week before sowing until just before sowing. Use fixed boom power sprayers set at 200-300 kPa and fixed with fan nozzles. Use 120-500 litres water per hectare. Openings in screens should be equal to or larger than 80 mesh. Agitate by mechanical or hydraulic means in the spray tank. When spraying avoid overlapping. Shut off spray booms while starting, turning, slowing and stopping otherwise crop injury may result.

CropSure Metribuzz 750WG Herbicide alone

If possible spray and incorporate within the same operation. If not possible incorporate spray within 24 hours of application using Go-devil or Lilliston cultivators set to work 5 cm deep.

CropSure Metribuzz 750WG Herbicide plus Trifluralin 480

After spraying mix soil within 4 hours. Spray and incorporate during the one operation if possible. (Refer to Trifluralin 480 label for specific incorporation details).

SPRAYER CLEANUP

Thoroughly clean all traces of CropSure Metribuzz 750WG Herbicide and /or trifluralin 480 from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

DO NOT apply under meteorological conditions or from spraying equipment, which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, croplands or pastures.

Tomatoes

DO NOT plant follow-on crops of cucurbits or capsicums when applying CropSure Metribuzz 750WG Herbicide to autumn plantings.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply (except as recommended for crop use) or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots, as injury or loss may occur.

DO NOT plant treated area to sensitive crops such as Brassicas, sunflowers, beetroot, cucurbits, lettuce or onions for at least 6 months following application of CropSure Metribuzz 750WG Herbicide as crop injury may result. Thoroughly work the soil prior to planting of these crops.

DO NOT apply under meteorological conditions or from spraying equipment, which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, croplands or pastures.

Tomatoes

DO NOT plant follow-on crops of cucurbits or capsicums when applying CropSure Metribuzz 750WG Herbicide to autumn plantings.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury containers in a local landfill. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) that can be obtained from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE