

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **CropSure**

# **Bullion 960**

## **HERBICIDE**

Active Constituent: 960g/L S-METOLACHLOR  
Solvent: 10g/L Polyethanoxy (15) tallow amine

GROUP	15	HERBICIDE
-------	----	-----------

Controls certain annual grasses and broadleaf weeds in certain crops as specified in the directions for use

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS: 20 – 1000 LITRES

**CropSure Pty Ltd**  
18 Raymond Road  
Laverton North  
Victoria 3026  
ABN: 17 643 829 190  
ACN: 643 829 190  
Phone: 03 9931 2200  
[www.cropsure.com](http://www.cropsure.com)

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Will damage eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length pvc gloves and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length pvc gloves and face shield or goggles. If product in eyes wash it out immediately with water. Wash hands after use after each day's use, wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

## CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

<b>IN EMERGENCY DIAL</b> <b>000</b> <b>POLICE OR FIRE BRIGADE</b>	<b>24HR EMERGENCY RESPONSE:</b> <b>AU +61 1800 951 288</b> <b>NZ +64 800 700 112</b>
---	--

APVMA APPROVAL NO: 93862 / 140725

BATCH NO:

DATE OF MANUFACTURE:

May cause an allergic skin reaction. Very toxic to aquatic life with long lasting effects. Avoid breathing fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves, protective clothing and eye or face protection. Wash contaminated clothing before reuse. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice. In case of fire: Use carbon dioxide, dry chemical, foam, to extinguish. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

**CAUTION  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **CropSure Bullion 960 HERBICIDE**

Active Constituent: 960g/L S-METOLACHLOR  
Solvent: 10g/L Polyethanoxy (15) tallow amine

GROUP	15	HERBICIDE
-------	----	-----------

Controls certain annual grasses and broadleaf weeds in certain crops as specified in the directions for use

**IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No.: 93862 / 140725**

**CropSure Pty Ltd**  
18 Raymond Road  
Laverton North  
Victoria 3026  
ABN: 17 643 829 190  
ACN: 643 829 190  
Phone: 03 9931 2200  
[www.cropsure.com](http://www.cropsure.com)

## DIRECTIONS FOR USE

### Restraints

DO NOT apply to waterlogged soils

DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 2 days of application

DO NOT irrigate to the point of run off for at least 2 days after application

### SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

Crops	Weeds Controlled	State	Rate	Critical Comments
<b>Broccoli, Brussels sprouts, Cabbages, Cauliflowers</b>	Barnyard Grass, Blackberry Nightshade, Chickweed (suppression), Deadnettle, Fat-hen, Lovegrass, Pigeon Grass, Pigweed (suppression), Potato Weed, Shepherds Purse, Stinging Nettle, Summer Grass, Wireweed (suppression)	Qld, NSW, Vic, Tas, NT only	1.5 to 2 L/ha	Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied with 24 hours. Use rates towards the higher end of the range where Blackberry Nightshade, Chickweed, Fat-hen, Lovegrass, Pigeon Grass, Pigweed, Potato Weed and Wireweed are the major problem.  <b>See General Instructions for Warning – Brassica Crops</b>
	Crowsfoot Grass, Fierce Thornapple (suppression), Liverseed Grass, Winter Grass	NSW only		Apply immediately after transplanting. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Apple of Peru, Common Sowthistle, Fierce Thornapple, Liverseed Grass, Redroot Amaranth, Twiggy Turnip and Nettleleaf Goosefoot are the major problems.
	Apple of Peru, Awnless Barnyard Grass, Fierce Thornapple (suppression) Liverseed Grass, Yellow Weed, Redroot Amaranth (suppression)	Qld only		<b>WARNING: Qld only:</b> Stunting and reduced yield may occur in broccoli in red soils and light alluvial soils.

	Annual Ryegrass, Common Cotula, Common Sowthistle (suppression), Crowsfoot Grass, Fiddle Dock, Italian Ryegrass, Liverseed Grass, Nettleleaf Goosefoot (suppression), Twiggy Turnip (suppression), Winter Grass.	Vic, Tas only		<b>See General Instructions for Warning – Brassica Crops</b>
<b>Canola</b>	Toad Rush	NSW, SA, Vic, Tas Qld	0.15 to 0.25 L/ha	<p>Apply at or immediately after planting and before crops and weeds emerge. Apply to moist soil, use rates towards the higher end of the range where longer residual control is required.</p> <p><b>See General Instructions for use in Zero Tillage systems</b></p>

Crops	Weeds Controlled	State	Rate	Critical Comments
<b>Cotton</b>	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew	Qld, NSW, WA only	1 L/ha	<p><b>Pre-emergent.</b> Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare.</p> <p>Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected.</p> <p>DO NOT throw untreated soil onto treated areas as this will reduce weed control.</p> <p>DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50% to 70%.</p> <p><b>See General Instructions for Warning – Cotton</b></p>
<b>Green Beans Navy Beans</b>	Black Berry Nightshade, Common Sowthistle, Fat-hen, Potato Weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass and suppression of Pigweed.	Qld, NSW, Vic, SA only	1.5 to 2 L/ha	<p><b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated.</p> <p>Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm.</p> <p>DO NOT throw untreated soil onto treated areas as this will reduce weed control.</p>

	Powell's Amaranth	Tas only	2 L/ha	<p><b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated</p> <p>For sufficient Amaranth control the CropSure Bullion 960 Herbicide application should be followed by 1 L/ha bentazone 480 (2 leaf Amaranth) plus 2L/ha bentazone 480 (4 leaf Amaranth). Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control.</p>
--	-------------------	----------	--------	---

Crops	Weeds Controlled	State	Rate	Critical Comments
<b>Maize Sweet Corn</b>	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew.	All States	1 to 2 L/ha	<p><b>Pre-emergent:</b> Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected.</p> <p>DO NOT throw untreated soil onto treated areas as this will reduce weed control.</p>
	Blackberry nightshade, Caltrop, Common Thornapple, Mintweed, Needle Burr, Noogoora Burr, Pigweed, Redroot Amaranth, Slim Amaranth		1 to 2 L plus 2.5 L of Flowable Atrazine (500g/L) /ha	<p>DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50% to 70%.</p>
<b>Pastures –</b> Trikkala Sub-clover, Haifa white clover, Maral Persian Clover, Paradana Balansa clover, Redquin Red Clover, Phalaris and Cocksfoot.	Toad Rush	NSW, Vic, Sa, Tas, WA, NT only	200 to 250 mL/ha	<p>Apply at or immediately after planting and before pasture species and weeds emerge. Application should be made to a moist seed bed. Sufficient rain to thoroughly wet the top 3 to 4cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required.</p> <p>DO NOT use in soils where prolonged waterlogging is anticipated otherwise unacceptable crop retardation will occur. Avoid damage to pasture species by avoiding overlaps and double spraying in headlands where White, Persian and</p>

				Balansa clovers are sown and if the 250mL rate is being used.
<b>Peanuts</b>	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrass, Pigeon Grass, Summer Grass, Wandering Jew.	Qld, NSW, WA, NT only	1 to 2 L/ha	Apply before pre-irrigation where soil crusting is likely. Apply before, at, or immediately after planting and before weeds germinate. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4 cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected.
<b>Soybeans Sunflowers</b>		All States		

Crops	Weeds Controlled	State	Rate	Critical Comments
<b>Sorghum treated with Concep II Sorghum Seed Safener</b>	Blackberry nightshade, Caltrop, Common Thornapple, Mintweed, Needle Burr, Noogoora Burr, Pigweed, Redroot Amaranth, Slim Amaranth	Qld, NSW, WA, NT, only	1 to 2L/ha 1 to 2L/ha plus 2.5L/ha of flowable Atrazine (500g/L)	<b>Pre-emergent:</b> Apply before, at or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60 to 120 litres of mixture per hectare. DO NOT apply to sorghum seed not previously treated with Concep II Sorghum Seed Safener. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3 to 4 cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3 to 4 cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use the lower rate on sandy soils.
<b>Sugar Cane plant and ratoon</b>	Awnless Barnyard Grass, Barnyard Grass, Bellvine, Billygoat Weed (Blue Top). Blackberry Nightshade, Caltrop, Common Sowthistle (Milk Thistle), Common Thornapple, Crowsfoot Grass, Green Summer Grass, Guinea Grass, Liverseed Grass, Mexican Clover, Mintweed, Needle Burr, Passionfruit Vines, Pigeon Grass, Pigweed, Redroot Amaranth, Square Weed, Star of Bethlehem, Summer Grass, Wild Rose	Southern Qld only (Bundaberg South)	1.1 to 1.45 L/ha plus 2.7 to 3.6 L/ha of flowable Atrazine (500g/L) or 1.5 to 2kg/ha of Atrazine (900g/kg) granules. (For example, CropSure Atrazine 900WG Herbicide)	Apply as a pre- or post-emergent application to crop. Apply once per year only. Application should be made to a moist soil before weeds and grasses have germinated. If conditions remain dry for a period of 10 days after spraying, irrigation or a shallow cultivation (2.5 cm) may assist results. DO NOT throw untreated soil onto treated areas as this will reduce weed control. In Southern Queensland use rates towards the higher end of the range on heavy soils where a high grass population is expected. In Northern Queensland application must be made to a moist soil and rainfall or irrigation should occur within 24 hours of application. Use rates towards the higher end of the range where high Green Summer Grass population is expected. Where broadleaf weeds and grasses have emerged and are in the 2 to 4 true leaf stage, knockdown herbicides such as ametryn (500g/L at 2L/ha), a flowable diuron (500g/L at 1L/ha or Paraquat/Diquat (1L/ha), applied as a post or directed spray, can be added. <b>Diuron Tank Mixes:</b> Read and follow all label directions including restraints, spray drift restraints, mandatory no spray zones, critical comments, withholding periods, regional use restrictions, and safety directions for the tank mix products.
		Northern Qld only (Mackay North)	1.45 to 1.8 L/ha plus 2.7 to 3.6 to 4.5L/ha of flowable Atrazine (500g/L) or 2 to 2.5 kg/ha of Atrazine (900g/kg) granules.	Where broadleaf weeds only have emerged and are at the 2 to 4 true leaf stage then a 500g/L 2, 4-D amine at 2 L/ha should be added. In all cases add a non-ionic surfactant. Follow all instructions and restriction on the Atrazine labels.

Crops	Weeds Controlled	State	Rate	Critical Comments
<b>Sweet Potatoes</b>	Black Pigweed, Redshank, Potato weed, Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Lovegrasses, Pigeon Grass, Summer Grass, Suppression of Evening Primrose	Qld, NSW, Vic, SA, NT only	1.5 L/ha	<b>Pre-emergent:</b> Apply immediately after transplanting before weeds have germinated. Sufficient irrigation to wet the soil through the weed zone should be applied within 24 hours. Further weed germination may occur following rehilling due to exposure of untreated soil.
<b>Tobacco</b> (transplanted in field)	Barnyard Grass, Lovegrass, Pigeon Grass, Summer Grass, and suppression of Common Thornapple ( <i>Detura stramonium</i> ) and Nightshades ( <i>Solanum sarachoides</i> and <i>S. gigrum</i> )	NSW, Vic only	1.5 L/ha	Apply by low volume boom spray prior to transplanting and before weeds and grasses germinate. The seed bed should be fine and free of large clods. Thoroughly incorporate within 7 days of spraying using a rotary hoe or similar equipment set to work to a depth of 7.5cm. DO NOT use disc cultivating implements. An even coverage of the herbicide is essential. The herbicide needs to be activated by moisture through to the top 3 to 4cm of soil at the time of application or within 4 days of incorporation. This can be achieved by applying to moist soil, irrigating soon after incorporation or irrigating immediately after transplanting. Avoid boom overlaps as double treatment may cause growth suppression. Temporary crop retardation may be observed when soil conditions remain dry after application or where Black Root Rot may occur.
<b>Barley, Oats</b>	Suppression of Annual Ryegrass	NSW, Vic, Tas, SA, WA, ACT only	375 to 500 mL/ha	Apply to moist soils immediately after planting and before crop and weeds emerge. Use the lower rate on light soils. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Damage may occur if crop is sown too shallow (less than 4 cm). Damage may occur on light sandy soil types after heavy rain. <b>See General Instructions for use in Zero tillage Systems.</b>

Crops	Weeds Controlled	State	Rate	Critical Comments
<b>Wheat, Barley, Oats, Triticale</b>	Toad Rush	NSW, Vic, Tas, SA, WA, ACT only	150 to 250mL per ha	Apply at or immediately after planting and before crop and weeds emerge. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3 to 4cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required or in situations where crop competition is minimal. When mixed with Triasulfuron 750g/L (for example CropSure Logjam 750WG Herbicide), apply pre-sowing and incorporate by sowing operation (for wheat only). <b>See General Instructions for use in Zero tillage Systems.</b>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD**

**Canola:** DO NOT GRAZE TREATED FIELDS FOR 10 WEEKS AFTER APPLICATION.

**Green beans:** DO NOT HARVEST BEANS FOR 8 WEEKS AFTER APPLICATION  
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 10 WEEKS AFTER APPLICATION.

**Pastures:** DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

**Sweet potatoes:** DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 23 WEEKS AFTER APPLICATION.

**Wheat, Barley, Oats, Triticale:** DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.

**HARVEST all other crops:** NOT REQUIRED WHEN USED AS DIRECTED.

**FORAGE all other crops (Except Sweet Potatoes):** DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

## General Instructions

### Uses

The product is a short residual pre-emergent herbicide for the control of many important annual grasses and broadleaf weeds in the crops nominated. The product does not control the important broadleaf weeds in maize, sweet corn and Concep II treated sorghum, but when mixed with Flowable Atrazine control of many important annual weeds and grasses is obtained without the problem of carryover of herbicides affecting succeeding crops. The product acts mainly through shoot (coleoptile) absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the top 3 to 4cm of soil, where it must be present at the time of weed germination.

### Application

**Ground application:** Apply in a minimum of 60 L/ha of water.  
DO NOT apply by aircraft.

### Mixing

Fill the spray tank to one quarter full. Then add CropSure Bullion 960 Herbicide to make up the final spray volume. Pour CropSure Bullion 960 Herbicide into the stream of oncoming water. Good agitation at the time CropSure Bullion 960 Herbicide is added to the tank will ensure good mixing.

**When Tank Mixing:** Wettable powder or water dispersible granule formulations should be added to the tank first, followed by suspension concentrates (flowables) water solution salts then CropSure Bullion 960 Herbicide or other emulsifiable concentrate formulations. Maintain thorough agitation during mixing and application. Agitate tank mixes vigorously if allowed to stand. **Note:** Tank mix spray solution should NOT be left standing in the vat overnight.

**WARNING – Cotton:** DO NOT use when planting into dry soil or soil with marginal moisture. Under dry planting conditions heavy rain or flood irrigation soon after planting, together with below average soil temperatures, can result in crop injury, due to the inability of the germinating plant being able to metabolise above normal amounts of the herbicide that may accumulate in the root zone as a result of accelerated capillary movement CropSure Bullion 960 Herbicide should not be used in situations where these conditions are likely to occur.

**WARNING – Brassica crops:** Crop retardation may occur where CropSure Bullion 960 Herbicide is used on soils that have a low organic matter content and which contain more than 60% of fine sand and silt. On these soil types, use rates of CropSure Bullion 960 Herbicide towards the lower end of the rate range and apply onto moist soil. Irrigation after application should be limited to no more than 25mm. DO NOT use more than once per year.

**Zero tillage (Canola, Barley, Oats, Wheat, Triticale):** When sowing using knife points and zero tillage, and leaving a seeding slot, DO NOT apply post-seeding unless harrows (or similar) have filled in the seeding slot before application. Alternatively apply CropSure Bullion 960 Herbicide immediately after seeding.

### Resistance Warning

GROUP	15	HERBICIDE
-------	----	-----------

CropSure Bullion 960 Herbicide is a member of the amides group of herbicides with diverse sites of action. For weed resistance management this is Group 15 herbicide. Some naturally-occurring weed

biotypes resistant to CropSure Bullion 960 Herbicide and other group 15 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CropSure Bullion 960 Herbicide or other group 15 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Bullion 960 Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a CropSure Pty Ltd representative.

## **PRECAUTIONS**

### **Re-entry Period**

DO NOT enter treated area without protective clothing until spray has dried.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS**

DO NOT apply under weather conditions, or from spray equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands and pastures

DO NOT plant crops other than broccoli, brussels sprouts, cabbages, cauliflowers, cotton, maize, peanuts, soybeans, sunflowers, sweet corn, tobacco, or Concep II treated sorghum within six months of application of CropSure Bullion 960 Herbicide

## **PROTECTION OF LIVESTOCK**

Do not graze or cut sweet potato leaves for stock food.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

TOXIC TO FISH. DO NOT contaminate streams, rivers or watercourse with the chemical or used containers.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto adjacent areas, particularly wetlands, water bodies or watercourses.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## **SAFETY DIRECTIONS**

Will damage eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length pvc gloves and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a

washable hat, elbow-length pvc gloves and face shield or goggles. If product in eyes wash it out immediately with water. Wash hands after use after each day's use, wash gloves, face shield or goggles and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## **SAFETY DATA SHEET (SDS)**

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

## **CONDITIONS OF SALE**

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.