

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure

Dialon 700WG

Fungicide

ACTIVE CONSTITUENT: 700 g/kg DITHIANON

GROUP	M9	FUNGICIDE
--------------	-----------	------------------

For the control of certain diseases of stone fruit, pome fruit and vines as per the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 1 kg - 15 kg

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilisers.

Single-rinse or shake remainder into spray tank. Do not dispose of undiluted chemicals on site.

Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Harmful if inhaled. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. Do NOT inhale dust. Repeated exposure may cause allergic disorders. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: Australia 13 11 26; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

In a Transport Emergency
Dial 000
Police or Fire Brigade

24HR Medical Emergency Information:
AUSTRALIA +61 1800 951 288
NEW ZEALAND +64 800 700 112

APVMA Approval No.: 90750/144538

Batch No:

Date of Manufacture:

Harmful if swallowed. Very toxic to aquatic life with long lasting effects. Avoid breathing dusts. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Brush off loose particles from skin. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. In case of fire: Use carbon dioxide, dry chemical, foam, to extinguish. Store locked up. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure

Dialon 700WG

Fungicide

ACTIVE CONSTITUENT: 700 g/kg DITHIANON

GROUP	M9	FUNGICIDE
--------------	-----------	------------------

For the control of certain diseases of stone fruit, pome fruit and vines as per the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA APPROVAL NO: 90750/144538

CropSure Pty Ltd
18 Raymond Road
Laverton North
Victoria 3026
ABN: 17 643 829 190
ACN: 643 829 190
Phone: 03 9931 2200
www.cropsure.com

DIRECTIONS FOR USE**RESTRAINTS:**

DO NOT apply if it is likely to rain before spray is dry.

DO NOT apply to wet crops.

DO NOT apply under slow drying conditions as an increase in fruit russet may occur.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift
DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

RATE					CRITICAL COMMENTS
CROP	DISEASE	STATE	RATE/100L	WHP	
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.					For all uses in this table apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Apples	Black spot, scab (<i>Venturia inaequalis</i>)	All States	18 g/100 L water	21 Days (harvest)	On apples intended for cold store Do not apply later than 42 days from harvest. This does not apply to apples intended for controlled Atmosphere storage. Do not apply during blossoming, in Qld only.
					Apply at 7 to 10 day intervals over the primary infection period from green tip to petal fall and continue at 10 to 14 day intervals thereafter while conditions permit infection. The product can also be used as a protectant addition to the DMI (Group 3) fungicides.
	Bitter rot (<i>Glomerella cingulata</i>)				Apply at 14 to 21 day intervals from petal fall to 21 days before harvest, using the shorter interval during periods of high humidity. Applications in the latter half of the season will assist in the control of post harvest development of bitter rot. A program of sprays for black spot control in apples will also provide control of bitter rot.
Pears	Black spot, scab (<i>Venturia pirina</i>)	Vic only	35 g or 50 g /100 L water	21 Days (harvest)	Apply from finger stage every 10 to 14 days while conditions permit infection. Use the higher rate when and where conditions are highly favourable for primary infection.
		NSW, Tas, SA, WA only	75 g/100 L water		
		Qld only	100 g/100 L water		
Grape Vines	Downy Mildew (<i>Plasmopara viticola</i>)	All States	25 g/100 L water (apply a minimum of 500 L/ha at shoots 10- 15cm length and a minimum of 1000 L/ha thereafter) or by aircraft at 550 g/ha	21 Days (harvest)	Apply when shoots are 10cm long, prior to infection occurring. While conditions permit infection, continue applications at 7 - 10 day intervals in wet weather, decreasing to 21 days in dry weather. To protect bunches, apply during flowering and 7 - 10 days later, then every 10 to 21 days as above.
			50 g/100 L water (apply a minimum of 500 L/ha at shoots 10- 15cm length and a minimum of 1000 L/ha thereafter) or by aircraft at 550 g/ha		Apply when shoots are 10cm long, prior to infection occurring. While conditions permit infection, continue applications at 10 day intervals in wet weather, decreasing to 21 days in dry weather. To protect bunches, apply during flowering and 2 weeks later, then every 10 to 21 days as above. Use the higher rate when and where conditions are highly favourable for primary infection.

Black spot (<i>Elsinoe ampelina</i>)	50 g/100 L water	Apply every 2 weeks from bud-burst to flowering.
Phomopsis cane and Leaf Blight (<i>Phomopsis viticola</i>)	50 to 75 g /100 L water	Use the higher rate for Phomopsis control when and where conditions are favourable for disease development.

CROP	DISEASE	STATE	RATE/100L	WHP	CRITICAL COMMENTS
Canning Peaches	Brown Rot (<i>Monilinia fructicola</i>)	NSW, Vic, Tas, SA only	100 g/100 L water plus a non-ionic wetting agent	1 Day (harvest)	Apply according to local recommendations or at budswell, full bloom, petal fall, shuck fall and at 3 weeks and 1 to 7 days before harvest.
		Qld only			Apply according to local recommendations or at budswell, full bloom, petal fall, shuck fall and at 4 weeks, 2 weeks and 1 to 3 days before harvest.
Apricots, Cherries, Nectarines, Peaches, Plums, Prunes	Brown Rot (<i>Monilinia fructicola</i>)	NSW, Vic, Tas, SA only	21 Days (harvest)		Apply according to local recommendations or at budswell, full bloom, petal fall, shuck fall and at 3 weeks before harvest. Within 21 days of harvest use another registered fungicide.
		Qld only			Apply according to local recommendations or at budswell, full bloom, petal fall, shuck fall and at 4 weeks before harvest. Within 21 days of harvest use another registered fungicide.
Apricots, Nectarines, Peaches	Freckle (<i>Venturia carpophila</i>)	All states	75 g/100 L water plus a non-ionic wetting agent		Apply according to local recommendations or at early bloom and shuck fall and then at monthly intervals until 21 days before harvest.
Nectarines, Peaches	Leaf Curl (<i>Taphrina deformans</i>)	All states			Apply at early budswell. the addition of a summer grade spraying oil will improve effectiveness.
	Rust (<i>Uromyces spp.</i>)		50 g/100 L water plus a non-ionic wetting agent		Apply according to local recommendations or at about monthly intervals from shuck fall until 21days before harvest.
Plums, Prunes	Rust (<i>Uromyces spp.</i>)				Apply according to local recommendations or at about monthly intervals from shuck fall until 21days before harvest.
All stone Fruit	Shot Hole (<i>Stigmina carpophila</i>), Scab/Peach Blight		150 g/100 L water plus a non-ionic wetting agent		Apply according to local recommendations leaf fall and early to mid blossoming.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIOD:

CANNING PEACHES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

FRUIT OTHER THAN CANNING PEACHES: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Mixing

With the agitation system operating, pour the measured volume of CropSure Dialon 700WG Fungicide slowly into the required volume of water in the spray vat. Mix thoroughly and then add any other compatible spray materials. Sprays containing CropSure Dialon 700WG Fungicide should be used within 3 hours of preparation and they should be agitated continuously during this period.

Wetting Agents

The addition of a non-ionic wetting agent, such as SST Vitiwet, is required when CropSure Dialon 700WG Fungicide is being applied to stone fruit.

Where a Wetting Agent is not required for CropSure Dialon 700WG Fungicide, one may be added if required for other pesticides, provided it is non-ionic.

APPLICATION

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which supplies water volumes less than those required to reach the point of run-off) and matched the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can be calculated in the following way.

Example only

1. Dilute spray volume as determined above: For example 1500 L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentrate factor in this example is: $3 \times$ (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$).
4. If the dilute label rate is 10 mL/100 L, then the concentrate rate becomes 3×10 , that is 30 mL/100 L of concentrate spray.

- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as crop grows.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

GRAPE VINES

Application by Ground Equipment

When applying as a dilute spray, apply sufficient volume to wet all leaf surfaces to the point of run-off. Apply as a medium to fine spray preferably using cone nozzles. Air-blast sprayers are recommended for application to vines with very dense foliage.

Application by Aircraft

Apply in a minimum of 20 litres of water per hectare. May be applied with hydraulic nozzles or rotary atomizers operated to produce fine to medium droplet size category. Avoid application in calm or very windy conditions or when the temperature and humidity cause rapid drying.

Compatibility

CropSure Dialon 700WG Fungicide is compatible with any one of the following products: Kingfisher Systemic Fungicide, Myclonil WG Fungicide, Bayfidan 250EC, Chlorpyrifos WP, omethoate 800, Gusathion 350, methomyl, penconazole 100 EC, Torque and wettable sulphur.

CropSure Dialon 700WG Fungicide should NOT be mixed with more than one other material, other than non-ionic wetting agent.

CropSure Dialon 700WG Fungicide is incompatible with Agridex, Tokuthion, Chlorpyrifos EC, Iprodione (Rovral) and winter or summer oils, except where mentioned under stone fruit. It is also incompatible with alkaline materials such as lime sulphur, lime and Bordeaux mixture. Under certain conditions it is incompatible with spraying oils and some other liquid formulations (see below).

Spraying Oils: CropSure Dialon 700WG Fungicide should only be applied with Winter or Summer oils at the dormant or bud burst stages. At other times it should not be applied with these materials or before or after oil cover sprays.

Liquid Formulations: CropSure Dialon 700WG Fungicide should not be applied to pome fruit, Santa Rosa or Wilson plums mixed with liquid formulations liable to cause ring spot.

Crop safety

If climatic conditions at the time of application result in slow drying of the spray deposit, some varieties of fruit may show minor skin russetting.

FUNGICIDE RESISTANCE WARNING

GROUP	M9	FUNGICIDE
-------	----	-----------

CropSure Dialon 700WG Fungicide is a member of the Multi-site Activity Group of fungicides. For fungicide resistance management, CropSure Dialon 700WG Fungicide is a Group M9 fungicide. Some naturally occurring individual fungi resistant to the product and other Group M9 fungicide may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if this fungicide is used repeatedly. These resistance fungi will not be controlled by CropSure Dialon 700WG Fungicide, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Dialon 700WG Fungicide to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or watercourses with the chemical or the used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilisers.

Single-rinse or shake remainder into spray tank. Do not dispose of undiluted chemicals on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Harmful if inhaled. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. Do NOT inhale dust. Repeated exposure may cause allergic disorders. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: Australia 13 11 26; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

In a Transport Emergency Dial 000 Police or Fire Brigade
--