

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CropSure Diuron 900WG HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg DIURON

GROUP 5 HERBICIDE

For the control of weeds in asparagus, bananas, cereals, cotton, lupins and sugar cane as specified in the directions for use table.

IMPORTANT: READ THIS LABEL BEFORE OPENING OR USING THIS PRODUCT

NET CONTENTS: 10 -15 KG

CropSure Pty Ltd

18 Raymond Road
Laverton North
Victoria 3026

ABN: 17 643 829 190

ACN: 643 829 190

Phone: 03 9931 2200

www.cropsure.com

DIRECTIONS FOR USE**RESTRAINTS:**

DO NOT use in water logged areas

DO NOT apply if greater than 50 mm of rainfall is expected within 3 days of application

DO NOT irrigate within 3 days of application

DO NOT apply to field where the slope exceeds 3%

DO NOT spot spray more than 5% of total farm areas

DO NOT apply more than once per calendar year

DO NOT apply by air

Cotton

DO NOT use in cotton UNLESS ALL irrigation tailwater and/or up to 25 mm of rainfall can be captured and held on farm.

Irrigation channels and drains

DO NOT use in irrigation channels and drains UNLESS ALL irrigation tailwater and rainfall can be captured and held on farm.

Desert Channels (bore drains)

DO APPLY between 1 December and 30 March

DO NOT open drains for 72 hours following treatment

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

Boom Sprayers

DO NOT apply by a boom sprayer unless droplets are not smaller than a COARSE spray droplet size category.

MANDATORY NO-SPRAY ZONES

DO NOT apply where there is non-target vegetation downwind from the application area and within the mandatory no-spray zones shown in the table below.

DO NOT apply where there are aquatic and wetland areas including aquaculture ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in the table below.

Situation	Rate g/ha	Downwind buffer zone	
		Aquatic	Terrestrial
Wheat, barley, triticale, cereal rye, oats (WA only)	278 - 555	30	60
Wheat, barley, triticale, and oats (WA only)	200 - 278	15	30
Wheat, barley, oats (NSW, Vic, ACT, SA only)	500	25	50
Wheat, barley (SA only)	711 - 978	50	100
Wheat, barley (NSW, Vic, ACT, SA only)	278	15	30
Summer fallows (SA only)	278	15	30
Cotton	1000 - 2000	100	200
Lupins (WA only)	1100	30	80
Bananas	278 - 500	25	50
Sugarcane	2000	100	200
Asparagus	2000	100	200

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Asparagus	Annual broadleaf weeds and some annual grasses	Vic only	1.5 – 1.9 kg	Apply in established beds after they have been "hilled up" and before emergence of spikes. This will give 10 to 14 weeks freedom from weeds. Post-emergence application may be made, but the spray should be directed so as to contact only the base of the stems. Use the higher rates on heavier soils. DO NOT use on seedling asparagus.
Bananas	Amaranthus, Barley Grass, Barnyard Grass, Capeweed, Cobbler's Pegs, Crowsfoot Grass, Fat Hen, Guinea grass, Mustard, Pigweed, Prince of Wales feather, Ryegrass, Sowthistle, Summer Grass, Wild Oats, Wild Radish, Wild Turnip	Qld, NSW, WA only	275 - 500g Plus 1.2 or 1.6 L/ha Paraquat 250 (for example CropSure Parashot 250 Herbicide)	Add to Paraquat 250 (for example CropSure Parashot 250 Herbicide) to enhance weed control under favourable growing conditions and in open sunny conditions. CropSure Diuron 900WG Herbicide at rates up to 500 g/ha can be blanket sprayed. Use a directed spray for higher rates of CropSure Diuron 900WG Herbicide. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5 cm high use a minimum of 250 L spray solution/ha, increasing to 350 L/ha for weeds up to 10 cm high. Use a minimum spray volume of 400 L/ha for weeds > 10 cm high and for dense mature weeds. Refer to Paraquat 250 Herbicide label and full directions. If Paraquat 250 Herbicide rate is less than the ratio 500 mL/100 L add 100mL non-ionic surfactant 600g/L or 60 mL of non-ionic surfactant 1000g/100L spray mix.
Cotton	Annual broadleaf weeds and some annual grasses	Qld, NSW only	1 – 1.9 kg	Pre-emergence: Apply at planting or within 7 days of planting and prior to cotton emergence. Seed should be planted at least 4-5 cm deep and soil compacted over seeds by a normal planter packer wheel. Use high rate on soils high in clay or organic matter content. DO NOT use on light, sandy soils as injury to crop may occur. DO NOT make a second pre-emergent application to the same field during the same crop year as injury to the crop may result. Apply in 300-450 L water/ha. Post emergence (lay-by): Use the higher rate on heavier soil types high in clay and/or organic matter. DO NOT use on light, sandy soils low in organic matter as injury to the cotton may result. Apply when cotton is at least 30 cm high, or immediately after the last cultivation. If emerged annual weeds (15cm or less in height) are present add 600g/L non-ionic surfactant to the spray solution at a rate of 250-500 mL per 1000L. Apply CropSure Diuron 900WG Herbicide as a directed spray to cover the area beneath cotton plants and between rows. Avoid spray contact or drift onto cotton plants as injury may result. DO NOT apply more than once per calendar year. DO NOT re-plant treated areas to any crop within one (1) year after last application as injury may result, except cotton, corn or sorghum (grain sorghum only) which may be planted in the spring following treatment. Apply in 300-500 L water/ha.
Lupins (on light soils only)	Annual ryegrass, Capeweed, Clovers, Medics, Doublegee, Wild Radish, Wild Turnip Suppression only: Brome Grass, Dock, Sorrel, Wild Oats and Wireweed	WA only	1.1 kg	Use through boomspray equipment only. Apply in a minimum of 50 L water/ha. Higher volumes of water will give more even application. Must be applied prior to crop emergence. DO NOT use on white or grey sands

Crop	Weeds Controlled	State	Rate/ha	Critical Comments	
Sugar Cane (Plant and ratoon crops)	Amaranthus, Barley Grass, Barnyard Grass, Capeweed, Cobbler's Pegs, Crowsfoot Grass, Fat Hen, Guinea grass, Mustard, Pigweed, Prince of Wales feather, Ryegrass, Sowthistle, Summer Grass, Wild Oats, Wild Radish, Wild Turnip	Qld, NSW only	1.9 kg	<p>Post-emergence of cane APPLY ONLY as directed band spray over a maximum of 60% of the crop area. DO NOT apply in the Wet Tropics DO NOT apply in the Burdekin between 1 January and 29 February DO NOT apply in the Mackay/Whitsunday between 1 December and 30 April DO NOT apply in Mary-Burnett between 1 November and 29 February DO NOT apply in NSW between 1 November and 30 April</p> <p>Soil surface must be moist at all times of application. Rain or irrigation is required within 10 days of application. If weeds at early post-emergence stage, addition of 600g/L non-ionic surfactant at 250 mL/100 L water will enhance activity, but direct spray below cane leaves.</p>	
			275 – 1.9 kg Plus 1.2 - 1.6 L Paraquat 250 Herbicide (for example CropSure Parashot 250 Herbicide)	<p>APPLY ONLY as directed band spray over a maximum of 60% of the crop area. DO NOT apply in the Wet Tropics DO NOT apply in the Burdekin between 1 January and 29 February DO NOT apply in the Mackay/Whitsunday between 1 December and 30 April DO NOT apply in Mary-Burnett between 1 November and 29 February DO NOT apply in NSW between 1 November and 30 April</p> <p>Add to Paraquat 250 Herbicide to enhance weed control under favourable growing conditions and in open sunny conditions. CropSure Diuron 900WG Herbicide at rates up to 500 g/ha can be blanket sprayed. Use a directed spray for higher rates CropSure Diuron 900WG Herbicide. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5 cm high use a minimum of 250L spray solution/ha, increase to 350 L/ha for weeds up to 10 cm high. Use a minimum spray volume of 400 L/ha for weeds > 10 cm high and for dense mature weed. Refer to Paraquat label for rates and full directions.</p>	
Summer fallows	Potato weed (<i>Heliotropium europaeum</i>)	SA only	275 g	Only effective when mixed with Paraquat 250 Herbicide At 1.6 L/ha (See Paraquat 250 Herbicide label)	
Wheat, Barley, Oats	Capeweed, Charlock, Deadnettle, Iceplant, Melilotus, Mustard, Poppy, Saffron Thistle, Sheepweed, Three Cornered Jack, Turnips, Wild Radish, Yellow Burrweed	NSW, ACT, Vic, SA only	500 g	<p>Apply when weeds are at the 2-4 leaf stage and soon after the crop is past the 2-5 leaf stage which is normally within 6 weeks of sowing.</p> <p>DO NOT spray later than the early tillering stage.</p>	
Wheat, Barley, Tricale, Oats	Capeweed, Doublegee, London Rocket, Mustard, Wild Radish, Wild Turnip, Wild Mustard	WA only	200 g plus 400mL MCPA 500	Apply when the crop has at least 3-5 leaves on the main stem. (Zadoks 13-15).	<p>Apply by boomspray. Weeds must be in the 2-5 leaf stage. Crop and weeds should be dry at the time of spraying.</p> <p>DO NOT spray if rain is imminent. DO NOT apply to undersown medics or undersown sub-clover. Apply in 40-70 L water/ha.</p>
			280 g plus 250 mL 2,4-D Amine 500	Apply when the crop has at least 4-5 leaves on the main stem. (Zadoks 14-15)	

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Wheat, Barley	Amsinckia (Yellow Burrweed), Capeweed, Deadnettle, Doublegee, (Three Cornered Jack), Mustards, White Ironweed, (Sheepweed), Wild Radish	NSW, ACT and Vic only	280 g plus 500 mL MCPCA 500	Apply by boomspray when the crop is at the 3-5 leaf stage (NSW, SA only), 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. DO NOT spray if rain is imminent. DO NOT apply to undersown medics. In NSW DO NOT use on Durati, Kite, or Shorten Wheat. Apply in 20 -100 L/ha
	Capeweed, Doublegee, (Three Cornered Jack), Mustards, Wild Radish	SA only	280 g plus 350 mL MCPCA 500	
Irrigation Channels	Annual and Perennial Grasses and Broadleaf Weeds	Qld, Vic, SA, Tas only	39 g to 78 kg	Use during the non-crop season and when channel is not in use. It is necessary to "fix" it into the soil by applying CropSure Diuron 900WG Herbicide prior to expected rainfall. The soil should be moist and preferably bare, i.e., after regrading or cleaning channels If 50-100 mm of rain has not fallen after application and it is necessary to use the irrigation channel, fill it with water and allow to stand for 72 hours. DO NOT allow water to run to waste. DO NOT use in irrigation channels or drains UNLESS ALL irrigation tail water and rainfall can be captured and held on farm.
		NSW, and Vic only	19.5 kg	
Drainage Ditches				Apply under the same conditions as for irrigation channels. Apply in 1000 – 2000 water/ha.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:
NOT REQUIRED WHEN USED AS DIRECTED**

GENERAL INSTRUCTIONS

CropSure Diuron 900WG Herbicide is a powerful weedkiller which remains active in the soil for a long time. Use only at recommended rates and read label directions carefully before using. The effects of this product on weeds are slow to appear and will not become apparent until it is carried into the root zone of the weeds or weed seeds by moisture. Results vary with soil type – lower rates are effective on lighter soils and higher rates on heavier soils.

This product is most effective and economical when applied to moist bare soil or soil carrying a small amount of vegetation.

Application should be made shortly before the period of main growth and best results will be achieved if rainfall or irrigation occurs within 3-4 days of application. Where additional "knockdown" is required, 600g/L non-ionic surfactant at the rate of 250 – 500 mL/100 L of water should be added. If existing herbage is excessive, it is advisable to mix Paraquat 250 Herbicide with CropSure Diuron 900WG Herbicide. This product is inactivated by absorption onto charcoal and other carboniferous materials.

Resistant weeds warning

GROUP	5	HERBICIDE
-------	---	-----------

CropSure Diuron 900WG Herbicide is a member of the urea group of herbicides. CropSure Diuron 900WG Herbicide has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management CropSure Diuron 900WG Herbicide is a Group 5 herbicide. Some naturally occurring weed biotypes resistant to CropSure Diuron 900WG Herbicide and other inhibitors of photosynthesis at photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CropSure Diuron 900WG Herbicide or other Group 5 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, CropSure Pty Ltd accepts no liability for any losses that may result from the failure of CropSure Diuron 900WG Herbicide to control resistant weeds.

Mixing

This product mixes readily with water. Steadily add the required quantity to the spray vat, which should be 50-75% full. The agitation system should be running during addition of the product and during the spraying operation. If left to settle, stir/agitate thoroughly before spraying. Avoid drawing accumulated particles into outlet pipe. DO NOT pre-mix as a paste.

Wetting Agent

Add 600 g/L non-ionic surfactant at 125 mL/100 L of spray mixture when applying to emerged weeds as a foliar spray.

Compatibility

CropSure Diuron 900WG Herbicide is compatible with the following: Paraquat 200 g/L and 250 g/L formulations.

Weed Control on Non-Crop Land

CropSure Diuron 900WG Herbicide is an effective herbicide for the residual control of many annual and perennial grasses and broadleaf weeds on non-crop land such as irrigation and drainage ditches where bare ground is desired. Established perennials such as Bindweed, Johnson Grass, Nutgrass, and Russian Knapweed are not satisfactorily controlled at rates recommended. Duration of control depends on the amount of chemical applied, soil type, rainfall and particular weed species.

Selective Use in crops

For selective use in crops, it is important to apply recommended rates correctly. CropSure Diuron 900WG Herbicide is most effective when applied to soil which has been well tilled (clod free) and is free of weeds and trash. Where not recommended for overall application, care should be taken to avoid spray drift onto fruit or foliage. Wetting agents should only be used where specified.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT use for soil sterilisation near or under trees or shrubs.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container in a dry, cool, well-ventilated locked room or place away from children, animals, foods, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. Single rinse bag before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Puncture or shred and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (e.g. phone Australia 13 11 26; New Zealand 0800 764 766 or a doctor).

SAFETY DATA SHEET

Additional Information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

CONDITIONS OF SALE

CropSure Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product, except for any provisions or rights under the Australian Consumer Law which cannot be excluded. The supply of this product is on the express condition that the purchaser does not rely on CropSure Pty Ltd skill of judgement in the purchasing or using the same and every person using this product does so at their own risk absolutely. The Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions. This product is designed to be only used according with the label directions.

IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	24HR EMERGENCY RESPONSE: AU +61 1800 951 288 NZ +64 800 700 112
---	--

APVMA APPROVAL NO: 94026 /141284

BATCH NO:

DATE OF MANUFACTURE:

Harmful if swallowed. Suspected of causing cancer. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe dusts. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Do not eat, drink or smoke when using this product. Get medical advice or attention if you feel unwell. Brush off loose particles from skin. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. If exposed or concerned: Get medical advice. In case of fire: Use carbon dioxide, dry chemical, foam, to extinguish. Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.